

MEDICAL POLICY

Medical Policy Title	Hip Arthroplasty
Policy Number	7.01.96
Current Effective Date	October 15, 2026
Next Review Date	June 2027

Our medical policies are guides to evaluate technologies or services for medical necessity. Criteria are established through the assessment of evidence based, peer-reviewed scientific literature, and national professional guidelines. Federal and state law(s), regulatory mandates and the member's subscriber contract language are considered first in the determination of a covered service.

(Link to [Product Disclaimer](#))

POLICY STATEMENT(S)

Partial Hip Replacement

I. Partial hip replacement is **medically necessary**, for **ANY** of the following conditions when **ALL** of the associated criteria have been met:

A. Femoral head/neck fracture, with

1. Imaging shows a fracture of the femoral head or femoral neck; **and**
2. Conservative management or surgical fixation is not considered a reasonable option;

B. Avascular necrosis (AVN), with

1. Imaging that shows AVN with collapse of the femoral head; **and**
2. Symptoms include **BOTH** of the following:
 - a. Function-limiting pain at short distances (e.g., walking less than one-quarter mile, limiting activity to two (2) city blocks, the equivalent to walking the length of a shopping mall) for at least three (3) months' duration; **and***

*Criteria exception: Three (3) months of function-limiting pain is not required when the medical record clearly documents why provider-directed non-surgical management is inappropriate.

- b. Loss of hip function that interferes with the ability to carry out age-appropriate activities of daily living or the demands of employment;

AND

3. Failure of at least three (3) months of provider-directed non-surgical management.

*Criteria exception: Provider-directed non-surgical management may be inappropriate. The medical record must clearly document why provider-directed non-surgical management is inappropriate (e.g., collapse of the femoral head, inflammatory arthritis, advanced dysplasia).

II. Partial hip replacement is **not medically necessary** for **ANY** other indication, or condition, or when **ANY** of the following are present:

A. Active local or systemic infection;

Medical Policy: Hip Arthroplasty

Policy Number: 7.01.96

Page: 2 of 10

- B. Vascular insufficiency, significant muscular atrophy of the leg, or neuromuscular disease severe enough to compromise implant stability or post-operative recovery;
- C. Charcot joint;
- D. Inflammatory arthritis affecting **both** the femoral head **and** the acetabulum.

Total Hip Replacement

III. Total hip replacement is **medically necessary**, for **ANY** of the following conditions when **ALL** of the associated criteria have been met:

- A. Femoral Head/Neck Fracture
 - 1. Imaging shows a fracture of the femoral head or femoral neck; **and**
 - 2. Conservative management or surgical fixation is not considered a reasonable option;
- B. Osteoarthritis, Avascular Necrosis (AVN), Inflammatory Arthritis
 - 1. Imaging shows **ANY** of the following findings:
 - a. Tönnis Grade 2 - 3 osteoarthritis;
 - b. Avascular necrosis with collapse of the femoral head; **or**
 - c. Inflammatory arthritis affecting **BOTH** the femoral head and the acetabulum with joint space narrowing;

AND

- 2. Symptoms include **BOTH** of the following:
 - a. Function-limiting pain at short distances (e.g., walking less than one-quarter mile, limiting activity to two city blocks, the equivalent to walking the length of a shopping mall) for at least three (3) months duration;* **and**

*Criteria exception: Three (3) months of function-limiting pain is not required when the medical record clearly documents why provider-directed non-surgical management is inappropriate;

- b. Loss of hip function which interferes with the ability to carry out age-appropriate activities of daily living or demands of employment;

AND

- 3. Failure of at least three (3) months of provider-directed non-surgical management;*

*Criteria exception: Three (3) months of provider-directed non-surgical management is not required when the medical record clearly documents why provider-directed non-surgical management is inappropriate.

IV. A total hip replacement is **not medically necessary** for **ANY** other indication, condition, or when **ANY** of the following are present:

- A. Active local or systemic infection;

Medical Policy: Hip Arthroplasty

Policy Number: 7.01.96

Page: 3 of 10

- B. Vascular insufficiency, significant muscular atrophy of the leg, or neuromuscular disease severe enough to compromise implant stability or post-operative recovery;
- C. Individuals undergoing dialysis.

Revision of Hip Replacement

- V. Revision of hip replacement (partial or total) is **medically necessary** for an individual who has previously undergone a partial or total hip replacement, when **ANY** of the following post-operative conditions are present:
 - A. Recurrent prosthetic dislocation/subluxation that is unresponsive to provider-directed non-surgical management;
 - B. Aseptic loosening;
 - C. Periprosthetic joint infection;
 - D. Periprosthetic fracture;
 - E. Instability of the implant (e.g., disassembly, modular neck failure);
 - F. Leg length discrepancy;
 - G. Osteolysis without eccentric wear (wear of elevated rim liner without wear superiorly);
 - H. Elevated serum metal levels as diagnosis for adverse local tissue reaction (ALTR) secondary to corrosion;
 - I. Unexplained, function-limiting pain at short distances (e.g., walking less than one-quarter mile, limiting activity to two (2) city blocks, the equivalent to walking the length of a shopping mall) for greater than six (6) months that is unresponsive to provider-directed, non-surgical management.
- VI. Revision of hip replacement is **not medically necessary** for any other indication or condition.

RELATED POLICIES

Not Applicable

POLICY GUIDELINE(S)

Not Applicable

DESCRIPTION

Total hip replacement is a surgical technique in which the femoral head and neck are removed, and the femoral canal (marrow space) is reamed out. The damaged hip joint is replaced with an artificial prosthesis composed of two or three different components: (1) the head that replaces the original femoral head; (2) the femoral component (a metal stem placed into the femur); and (3) the acetabular component, which is implanted into the acetabulum. The stem may be secured using bone cement or press-fit for the bone to grow into it.

Medical Policy: Hip Arthroplasty

Policy Number: 7.01.96

Page: 4 of 10

The Tonnis Classification System is commonly used to describe the presence of osteoarthritis in the hips, with grading as follows:

- Grade 0: No signs of osteoarthritis.
- Grade 1: Sclerosis of the joint, with slight joint space narrowing and osteophyte formation and no or slight loss of femoral head sphericity.
- Grade 2: Small cysts in the femoral head or acetabulum, with moderate joint space narrowing and moderate loss of femoral head sphericity.
- Grade 3: Large cysts in the femoral head or acetabulum, severe joint space narrowing or obliteration of the joint space, and severe deformity and loss of sphericity of the femoral head.

Revision of hip replacement (partial or total) involves surgical reconstruction or replacement due to failure or complications of previous hip replacement.

Non-surgical management regarding the treatment of hip osteoarthritis is defined as any provider-directed, non-surgical treatment that has been demonstrated in the scientific literature to be efficacious and/or is considered reasonable care in the treatment of hip pain from osteoarthritis. The types of treatment can include but are not limited to relative rest/activity modification, weight loss, supervised physiotherapy modalities and therapeutic exercises, oral prescription and non-prescription medications, assistive devices (e.g., cane, crutches, walker, wheelchair), and/or intra-articular (i.e., steroid) injections.

SUPPORTIVE LITERATURE

In a meta-analysis, Smith and colleagues (2010) compared the clinical and radiological outcomes and complication rates of hip resurfacing (HRS) and total hip arthroplasty (THA). A systematic review was undertaken of all published (Medline, CINAHL, AMED, EMBASE) and unpublished or gray literature research databases up to January 2010. Clinical and radiological outcomes, as well as complications of HRS, were compared to those of THA, using risk ratio, mean difference, and standardized mean difference statistics. Studies were critically appraised using the CASP appraisal tool. A total of 46 studies were identified from 1,124 citations. These included 3,799 HRSs and 3,282 THAs. On meta-analysis, functional outcomes for subjects following HRS were better than or the same as for subjects with a THA, but there were statistically significant increases in incidence of heterotopic ossification, aseptic loosening, and revision surgery with HRS, compared to THA. The evidence base showed a number of methodological inadequacies, such as the limited use of power calculations and poor or absent blinding of both patients and assessors, possibly giving rise to assessor bias. The authors concluded that, on the basis of the current evidence base, HRS may have better functional outcomes than THA, but the increased risks of heterotopic ossification, aseptic loosening, and revision surgery following HRS indicate that THA is superior in terms of implant survival.

In a 2019 retrospective cohort study, Inoue et al compared post-operative complications and survivorship of total hip and knee arthroplasty in dialysis and renal transplantation patients. They included a total of 107 patients undergoing primary total joint arthroplasty (TJA), including 50 who were receiving dialysis and 57 who had a prior renal transplantation. The end point was defined as revision surgery secondary to post-operative complications. Researchers found a significantly higher

Medical Policy: Hip Arthroplasty

Policy Number: 7.01.96

Page: 5 of 10

rate of post-operative complications in the dialysis cohort (28%, n=14 of 50 joints) compared to the renal transplant cohort (7.1%, n= 4 of 57 joints). There was a higher rate of surgical site infection (SSI) and periprosthetic joint infection (PJI) in dialysis patients, compared with renal transplantation patients (18% versus 3.5%, $P=0.02$). In addition, there was an increased rate of revision surgery in the dialysis cohort, compared to transplant cohort (24% versus 3.5%, $P=0.002$). A multi-variate analysis considering demographics and comorbidities revealed that patients with renal transplantation were less likely to have revision surgery, compared to patients on dialysis at the time of arthroplasty (95 % CI, $P=0.031$), and demonstrated a strong trend for lower complications (95% CI, $P=0.76$), although the latter was not statistically significant. Researchers concluded that transplantation was independently associated with reduced rates of revision surgery in the setting of chronic renal failure, suggesting that those who are candidates may benefit from renal transplantation before undergoing elective TJA.

Chou et al (2021) compared outcomes of TJA, including THA and total knee arthroplasty, in individuals with end-stage renal disease who were either receiving dialysis or had undergone a kidney transplant (KT). The analysis included 22 studies encompassing 9,384 patients (n = 8,921 dialysis; n = 463 KT). The overall mortality rate after TJA was 14.9%-13.8% in dialysis patients and 15.8% in KT patients. The overall surgical site complication (SSC) rate was 3.3%, with KT patients experiencing a rate of 3.6%. SSCs included hematoma, seroma, delayed wound healing, and superficial infections. The overall PJI rate was 3.9%, with dialysis patients at 4.0% and KT patients at 3.7%. Multivariate regression analysis found no statistically significant differences in outcomes between dialysis and KT patients. Both groups demonstrated elevated risks compared to the general population, with comparable outcomes between them.

Kim et al (2022) conducted a systematic review and meta-analysis of 10 retrospective studies that included 3,631,861 patients and 14,996 solid organ transplantations (SOT). They evaluate whether patients with a history of SOT experienced different outcomes after primary hip arthroplasty compared to non- SOT patients. Medical complications included cardiac complications, pneumonia, acute kidney injury, and pulmonary thromboembolism and deep vein thrombosis. The study showed that patients with a history of SOT undergoing hip arthroplasty are at higher risk for medical complications, transfusion, readmission, and short-term mortality compared to non-transplant patients. However, surgical complication rates such as infection and revision are not significantly different.

PROFESSIONAL GUIDELINE(S)

In the 2023 American Academy of Orthopaedic Surgeons adopted the evidence-based Clinical Practice Guidelines for the Management of Osteoarthritis of the Hip that state:

- “They strongly recommend conservative measures such as physical therapy, corticosteroid or hyaluronic acid injections, pharmacological management (NSAIDs).”

In the 2008 International Cartilage Repair Society Guidelines for the Management of Hip and Knee Osteoarthritis (OA), Part II, stated:

- “Patients with hip or knee OA who are not obtaining adequate pain relief and functional

Medical Policy: Hip Arthroplasty

Policy Number: 7.01.96

Page: 6 of 10

improvement from a combination of non-pharmacological and pharmacological treatment should be considered for joint replacement surgery. Replacement arthroplasties are effective, and cost-effective interventions for patients with significant symptoms, and/or functional limitations associated with a reduced health-related quality of life, despite conservative therapy.” (Zhang et al 2008)

REGULATORY STATUS

Not Applicable

CODE(S)

- Codes may not be covered under all circumstances.
- Code list may not be all inclusive (AMA and CMS code updates may occur more frequently than policy updates).
- (E/I)=Experimental/Investigational
- (NMN)=Not medically necessary/appropriate

CPT Codes

Code	Description
27125	Hemiarthroplasty, hip, partial (e.g., femoral stem prosthesis, bipolar arthroplasty)
27130	Arthroplasty, acetabular and proximal femoral prosthetic replacement (total hip arthroplasty), with or without autograft or allograft
27132	Conversion of previous hip surgery to total hip arthroplasty, with or without autograft or allograft
27134	Revision of total hip arthroplasty; both components, with or without autograft or allograft
27137	Revision of total hip arthroplasty; acetabular component only, with or without autograft or allograft
27138	Revision of total hip arthroplasty; femoral component only, with or without allograft

Copyright © 2026 American Medical Association, Chicago, IL

HCPCS Codes

Code	Description
Not Applicable	

ICD10 Codes

Medical Policy: Hip Arthroplasty

Policy Number: 7.01.96

Page: 7 of 10

Code	Description
M05.051-M08.959	Inflammatory polyarthropathies (hip) (code range)
M12.551 - M12.559	Traumatic arthropathy, hip (code range)
M16.0 - M16.9	Osteoarthritis of hip (code range)
M80.051A -M80.059S, M80.851A- M80.859S, M84.451A-M84.453S, M84.459A-M84.459S, M84.551A-M84.559S, M84.651A-M84.659S	Pathologic fracture of neck of femur (hip) (code range)
M84.750A-M84.759S	Atypical femoral fracture (code range)
M87.051-M87.059, M87.151-M87.159, M87.251-M87.256, M87.351-M87.353, M87.851-M87.859, M90.551-M90.559	Osteonecrosis of femur and thigh (code range)
M97.01XA-M97.02XS	Periprosthetic fracture around internal prosthetic hip joint (code range)
S72.001A-S72.26XS	Fracture of head and neck of femur (code range)

REFERENCES

Adrados M, et al. Institutional adherence to the American Association of Hip and Knee Surgeons body mass index guidelines lowers perioperative emergency department visits in primary total knee arthroplasty. The Journal of Arthroplasty. 2023 Jun;38(6S):S88-S93.

American Academy of Orthopaedic Surgeons (AAOS) [Internet]. Appropriate use criteria: Osteoarthritis of the hip: management. Dec 2017. [accessed 2026 May 11]. Available from: <https://www.orthoguidelines.org/go/auc/>

American Academy of Orthopaedic Surgeons (AAOS) [Internet]. Evidence-based clinical practice guideline: Management of osteoarthritis of the hip. Dec 2023. [accessed 2026 May 11]. Available from: <https://www.aaos.org/globalassets/quality-and-practice-resources/osteoarthritis-of-the-hip/oah-cpg.pdf>

Bannuru RR, et al. OARSI guidelines for the non-surgical management of knee, hip, and polyarticular

Medical Policy: Hip Arthroplasty

Policy Number: 7.01.96

Page: 8 of 10

osteoarthritis. Osteoarthritis Cartilage. 2019 Nov;27(11):1578-1589.

Bartels S, et al. Total hip arthroplasty leads to better results after low-energy displaced femoral neck fracture in patients aged 55 to 70 years: a randomized controlled multicenter trial comparing internal fixation and total hip arthroplasty Journal of Bone Joint Surgery. 2022;Aug;104(15):1341-1351.

Bhandari M, et al. Total hip arthroplasty or hemiarthroplasty for hip fracture. N Engl J Med. 2019 Dec 5;381(23):2199-2208.

Blankstein M, et al. What factors increase revision surgery risk when treating displaced femoral neck fractures with arthroplasty: a secondary analysis of the HEALTH Trial. J Orthop Trauma. 2020 Nov;34 Suppl 3:S49-S54.

Browne J, et al. Peritoneal dialysis does not carry the same risk as hemodialysis in patients undergoing hip or knee arthroplasty. J Bone Joint Surg. 2019;101(14):1271-77.

Carlo JO, et al. Perioperative implications of end-stage renal disease in orthopaedic surgery. J Am Acad Orthop Surg. 2015 Feb;23(2):107-18.

Cavanaugh PK, et al. Complications and mortality in chronic renal failure patients undergoing total joint arthroplasty: A comparison between dialysis and renal transplant patients. J Arthroplasty. 2016 Feb;31(2):465-72.

Chalmers BP, et al. Primary total hip arthroplasty for charcot arthropathy is associated with high complications but improved clinical outcomes. J Arthroplasty. 2018 Sept;33(9):2912-2918.

Chammout G, et al. HOPE-Trial: hemiarthroplasty compared with total hip arthroplasty for displaced femoral neck fractures in octogenarians: a randomized controlled trial. JB JS Open Access. 2019 May 1;4(2):e0059.

Chen W, et al. Direct anterior versus posterolateral approaches for clinical outcomes after total hip arthroplasty: a systematic review and meta-analysis. J Orthop Surg Res. 2020 Jun 23;15(1):231.

Chou TA, et al. Dialysis patients have comparable results to patients who have received kidney transplant after total joint arthroplasty: a systematic review and meta-analysis. EFORT Open Rev. 2021 Aug 10;6(8):618-628.

Douglas SJ, et al. Comparing Primary Total Hip Arthroplasty in Renal Transplant Recipients to Patients on Dialysis for End-Stage Renal Disease: A Nationally Matched Analysis. J Bone Joint Surgery Am. 2021; 103:2215-20.

Ekhtiari S, et al. Total hip arthroplasty versus hemiarthroplasty for displaced femoral neck fracture: a systematic review and meta-analysis of randomized controlled trials. J Bone Joint Surg Am. 2020 Sep 16;102(18):1638-1645.

Inoue D, et al. Comparison of postoperative complications and survivorship of total hip and knee arthroplasty in dialysis and renal transplantation patients. J Arthroplasty. 2020;35(4):971-75.

Inoue D, et al. Outcomes of simultaneous bilateral total hip arthroplasty for 256 selected patients in a single surgeon's practice. Bone Joint J. 2021 Jul;103-B(7 suppl B): 116-121.

Iorio R, et al. [Internet]. What are the absolute and relative contraindications to elective primary total

Medical Policy: Hip Arthroplasty

Policy Number: 7.01.96

Page: 9 of 10

joint arthroplasty (TJA), with respect to surgical site infection (SSI) and periprosthetic joint infection (PJI) risk? [Internet]. Second international consensus meeting on musculoskeletal infection 2018: hip and knee. [accessed 2026 May 11]. Available from:

https://www.icmortho.org/_files/ugd/0540e9_f047ffe9ecc543279bba6b84688bc27e.pdf

Jo S, et al. Clinical outcomes of total hip arthroplasty for displaced femoral neck fractures in patients 80 years of age and older selected by clinical frailty score. *Hip Pelvis*. 2020 Sep;32(3):148-155.

Kim CH, et al. Clinical outcomes following primary hip replacement arthroplasties in patients with solid organ transplantation: A systematic review and meta-analysis. *Hip Pelvis*. 2022 Sep;34(3):127-139.

Li J, et al. Comparison of total joint arthroplasty outcomes between renal transplant patients and dialysis patients-a meta-analysis and systematic review. *J Orthop Surg Res*. 2020 Dec 9;15(1):590.

Lieu D, et al. Total hip replacement in haemodialysis or renal transplant patients. *J Orthop Surg (Hong Kong)*. 2014 Dec;22(3):393-8.

Liu Y, et al. Comparing total hip arthroplasty and hemiarthroplasty for the treatment of displaced femoral neck fracture in the active elderly over 75 years old: a systematic review and meta-analysis of randomized control trials. *J Orthop Surg Res*. 2020 Jun 11;15(1):215.

Martin SD, et al. Hip arthroscopy versus physical therapy for the treatment of symptomatic acetabular labral tears in patients older than 40 Years: a randomized controlled trial. *The American Journal of Sport Medicine*. 2021 APR;49(5):1199-1208.

Popat R, et al. Outcomes of total hip arthroplasty in haemodialysis and renal transplant patients: systematic review. *Hip Int*. 2021 Mar;31(2):207-214.

Shrader MW, et al. Total hip arthroplasty in patients with renal failure: a comparison between transplant and dialysis patients. *J Arthroplasty*. 2006;21(3):324-329.

Smith TO, et al. The clinical and radiological outcomes of hip resurfacing versus total hip arthroplasty: a meta-analysis and systematic review. *Acta Orthop*. 2010;81(6):684-695.

Zhang W, et al. OARSI recommendations for the management of hip and knee osteoarthritis, part I: critical appraisal of existing treatment guidelines and systematic review of current research evidence. *Osteoarthritis Cartilage*. 2007 Sep;15(9):981-1000.

Zhang W, et al. OARSI recommendations for the management of hip and knee osteoarthritis, part II: OARSI evidence-based, expert consensus guidelines. *Osteoarthritis Cartilage*. 2008 Feb;16(2):137-62.

Zhang W, et al. OARSI recommendations for the management of hip and knee osteoarthritis, part III: Changes in evidence following systematic cumulative update of research published through January 2009. *Osteoarthritis Cartilage* 2010;18(4):476-499.

SEARCH TERMS

Not Applicable

CENTERS FOR MEDICARE AND MEDICAID SERVICES (CMS)

Medical Policy: Hip Arthroplasty

Policy Number: 7.01.96

Page: 10 of 10

[Total Joint Arthroplasty \(LCD L36039\)](#) [accessed 2026 May 11].

[Total Joint Arthroplasty \(LCA A57428\)](#) [accessed 2026 May 11].

PRODUCT DISCLAIMER

- Services are contract dependent; if a product does not cover a service, medical policy criteria do not apply.
- If a commercial product (including an Essential Plan or Child Health Plus product) covers a specific service, medical policy criteria apply to the benefit.
- If a Medicaid product covers a specific service, and there are no New York State Medicaid guidelines (eMedNY) criteria, medical policy criteria apply to the benefit.
- If a Medicare product (including Medicare HMO-Dual Special Needs Program (DSNP) product) covers a specific service, and there is no national or local Medicare coverage decision for the service, medical policy criteria apply to the benefit.
- If a Medicare HMO-Dual Special Needs Program (DSNP) product DOES NOT cover a specific service, please refer to the Medicaid Product coverage line.

POLICY HISTORY/REVISION

Committee Approval Dates

12/20/18, 06/20/19, 12/19/19, 12/17/20, 04/15/21, 04/21/22, 04/20/23, 10/17/24, 06/26/25, 10/16/25, 06/18/26

Date	Summary of Changes
06/18/26	• Annual review; policy intent unchanged.
10/16/25	• Off cycle review. Removed “and on a renal transplant list” from the not medically necessary indications for total hip.
06/26/25	• Annual review, policy statement regarding bilateral simultaneous hip replacement was removed from the policy.
01/01/25	• Summary of changes tracking implemented.
06/21/18	• Original effective date