

Pharmacy Management Drug Policy

SUBJECT: Ophthalmic Vascular Endothelial Growth Factor (VEGF) Inhibitors		
POLICY NUMBER: PHARMACY-138		
EFFECTIVE DATE: 04/2026		
LAST REVIEW DATE: 08/07/2026		
<i>If the member's subscriber contract excludes coverage for a specific service or prescription drug, it is not covered under that contract. In such cases, medical or drug policy criteria are not applied. This drug policy applies to the following line/s of business:</i>		
Policy Application		
Category:	☒ Commercial Group (e.g., EPO, HMO, POS, PPO)	☒ Medicare Advantage
	☒ On Exchange Qualified Health Plans (QHP)	☐ Medicare Part D
	☒ Off Exchange Direct Pay	☒ Essential Plan (EP)
	☒ Medicaid & Health and Recovery Plans (MMC/HARP)	☒ Child Health Plus (CHP)
	☐ Federal Employee Program (FEP)	☐ Ancillary Services
	☒ Dual Eligible Special Needs Plan (D-SNP)	

DESCRIPTION:

Vascular endothelial growth factor (VEGF) is a protein produced by cells in the body which stimulates the growth of new blood vessels. Occasionally cells can produce too much VEGF causing abnormal new blood vessels to grow in the eye (neovascularization). Abnormal neovascularization can cause leakage of blood and fluids into the retina, scar the macula, and affect vision. Ophthalmic vascular endothelial growth factor (VEGF) inhibitors block the action of VEGF which reduces leaking abnormal blood vessels in the retina. This slows or stops damage from the abnormal blood vessels and slows down vision loss. In some instances, VEGF-inhibitors may also improve vision. Ophthalmic VEGF inhibitors are administered as intravitreal injections. Ophthalmic conditions treated with VEGF inhibitors include neovascular (wet) age-related macular degeneration (nAMD), diabetic retinopathy (DR), diabetic edema (DME), macular edema following retinal vein occlusion (RVO), myopic choroidal neovascularization (mCNV) and retinopathy of prematurity (ROP).

Neovascular (wet) Age-Related Macular Degeneration (nAMD)

Neovascular Age-Related Macular Degeneration is a chronic, progressive disorder of the retina that is characterized by choroidal neovascularization (CNV), which is caused due to over-expression of VEGF.¹ When VEGF levels rise, they promote growth of abnormal vessels, vascular leakage, and hemorrhage underneath the macula, which cause the central vision to appear distorted, and if left untreated may lead to irreversible loss of vision.¹ Early stages of the disease are characterized by symptoms such as central vision blurring, difficulty with fine visual tasks, and metamorphopsia, defined as a type of visual distortion that causes linear objects to appear rounded.² Without medical intervention, nAMD progresses rapidly and is currently a leading cause of blindness in the older adult population in developed countries.² Introduction of intravitreal VEGF inhibitors has altered the natural disease course, reducing the rates of severe vision loss and improving outcomes of visual acuity in many patients.³ According to the American Academy of Ophthalmology (AAO), anti-VEGF therapy is currently considered first line therapy for nAMD.¹

Diabetic Retinopathy (DR) and Diabetic Macular Edema (DME)

Diabetic Retinopathy is a complication of chronic hyperglycemia, which causes capillary dropout, ischemia, inflammation, and ultimately upregulation of VEGF.⁴ Diabetic Macular Edema occurs due to breakdown of the blood-retina barrier and fluid accumulation inside the macula. While DME is a major cause of vision impairment in patients suffering from diabetes, randomized controlled trials indicated that therapy with intravitreal VEGF inhibitors was more effective in improving visual acuity, when

Pharmacy Management Drug Policy

Ophthalmic Vascular Endothelial Growth Factor (VEGF) Inhibitors

compared to laser photocoagulation.⁵ Both the American Diabetes Association (ADA) and AAO guidelines recommend anti-VEGF therapy as first-line for the treatment of center-involved DME and vision-threatening DR.⁴

Macular Edema Following Retinal Vein Occlusion (RVO)

Types of RVO include branch RVO and central RVO. Both are known to cause retinal ischemia, cystoid macular edema, and elevation of VEGF.⁶ The most common patient presentation is a sudden, pain-free loss of vision. VEGF inhibitors are the current standard of care for macular edema following RVO, with randomized clinical trials displaying improvement in visual acuity and anatomic outcomes, as a result of VEGF inhibition.⁶ Anti-VEGF therapy is preferred over corticosteroid, due to its superior efficacy and safety in patients suffering from RVO-associated macular edema.⁶

Myopic Choroidal Neovascularization (mCNV)

Patients with pathologic myopia are predisposed to CNV similar to that seen in patients with AMD.⁷ Therapy with VEGF inhibitors has demonstrated superior efficacy when compared to photodynamic therapy in visual acuity and resolution of active lesions.⁷

Retinopathy of Prematurity (ROP)

In premature infants, incorrect development of retinal vascularity leads to a condition called retinopathy of prematurity, in which dysregulation of VEGF leads to neovascularization and detachment of the tractional retina. Currently, aflibercept and bevacizumab are used in specialized centers as adjunct therapy, or in cases where laser therapy is contraindicated.^{8,9,10}

Ophthalmic Anti-VEGF Agents

- **Aflibercept (Eylea[®], Eylea HD[™], biosimilars to Eylea, Pavblu[™], Yesafili[™]):** approved for the treatment of nAMD, DME, DR, RVO, ROP^a.
- **Bevacizumab (Avastin[®], biosimilars):** compendial-approved for the treatment of nAMD, DME, DR, RVO and ROP.
- **Brolucizumab (Beovu[®]):** approved for the treatment of nAMD and DME.
- **Faricimab (Vabysmo[®]):** approved for the treatment of nAMD, DME and RVO.
- **Ranibizumab (Lucentis[®], Byooviz[®], Cimerli, Nufymco, Susvimo[™]):** approved for the treatment of nAMD, DME, DR, RVO, and mCNV^b.
 - a. Eylea only
 - b. Lucentis, Byooviz, Cimerli and Nufymco only

Comparative Safety and Efficacy

Large comparative trials, such as CATT, VIEW, HARBOR, DRDR.net Protocol T, and TENAYA/LUCERNE, have demonstrated similar improvements in visual acuity across available agents, assuming they are being administered at the recommended dosing intervals.^{3,4,5,6} Differences between products appear to include flexibility of dosing interval, drying effects and safety concerns, but not significant superiority in vision outcomes. Class-wide adverse events include endophthalmitis, intraocular inflammation, retinal detachment, and increases in intraocular pressure, which appears to be transient.¹¹ Although rare thromboembolic events have been observed, there is still no clear causal association between the medication class and this adverse event.¹¹ Drug-specific concerns are presented with Beovu and Susvimo, with the former being associated with retinal vasculitis and occlusive vasculopathy and the latter with implant-specific complications (dislocation, conjunctival erosion, increased risk of infection).^{15,16}

Pharmacy Management Drug Policy

Ophthalmic Vascular Endothelial Growth Factor (VEGF) Inhibitors

POLICY:

	Neovascular (Wet) Age-Related Macular Degeneration (nAMD)	Diabetic Macular Edema (DME)	Diabetic Retinopathy (DR)	Macular Edema following Retinal Vein Occlusion (RVO)	Myopic Choroidal Neovascularization (mCNV)	Retinopathy of Prematurity (ROP)
Beovu* (brolucizumab)	X	X				
Eylea* (aflibercept)	X	X	X	X		X
Pavblu* (aflibercept-ayyh)	X	X	X	X		
Yesafili* (aflibercept-ibvfv)	X	X	X	X		
Eylea HD* (aflibercept)	X	X	X	X		
Byooviz* (ranibizumab-nuna)	X	X	X	X	X	
Cimerli* (ranibizumab-eqrm)	X	X	X	X	X	
Lucentis* (ranibizumab)	X	X	X	X	X	
Nufymco* (ranibizumab-leyk)	X	X	X	X	X	
Susvimo* (ranibizumab)	X	X	X			
Vabysmo* (faricimab-svoa)	X	X		X		

*designated as a preferred product; however, preferred status still requires prior use of a bevacizumab-containing product unless otherwise specified

- The prescribed medication must be used for the indication(s) consistent with the population outlined in the table above **AND**
- Must be prescribed by an ophthalmologist **AND**
- Must be 18 years of age or older
 - Does not apply to Eylea for the diagnosis of Retinopathy of Prematurity (ROP) **AND**
- For a diagnosis of Neovascular (Wet) Age-Related Macular Degeneration (nAMD), Diabetic Macular Edema (DME), Diabetic Retinopathy (DR), Macular Edema following Retinal Vein Occlusion (RVO), and Myopic Choroidal Neovascularization (mCNV)
 - Patients new to treatment must have had serious side effects or drug failure (defined as at least 3 injections that results in a suboptimal clinical response) of a bevacizumab-containing product
 - For Susvimo
 - For patients that achieved a positive clinical response to at least two injections of bevacizumab but subsequently experienced disease progression or loss of response, this requirement is satisfied.
 - For patients that did not achieve a positive clinical response to at least two injections of bevacizumab, documentation must be provided confirming that the patient previously responded to at least two injections of an alternative VEGF inhibitor (e.g., aflibercept, ranibizumab, brolucizumab)
 - For Eylea, Pavblu, Yesafili, and Eylea HD
 - For patients with a diagnosis of DME or DR with a baseline visual acuity score of 20/50 or worse, a trial of a bevacizumab-containing product will not be required **AND**
- Refer to individual prescribing information for approved dosing
- Eylea HD will not be covered for any non-FDA approved diagnoses (i.e., Retinopathy of Prematurity [ROP])

Pharmacy Management Drug Policy

Ophthalmic Vascular Endothelial Growth Factor (VEGF) Inhibitors

7. Approval Timeframe

Line of Business	Medical Initial	Medical Recert
SafetyNet (Medicaid, HARP, CHP, Essential Plan)	6 months	6 months
Commercial / Exchange	6 months	6 months
Medicare	2 years	2 years

POLICY GUIDELINES:

- Utilization Management is contract dependent. Refer to specific contract/benefit language for exclusions.
 - Coverage criteria may be dependent on the contract renewal date.
 - Coverage of drugs listed in this policy are contract dependent.
 - Not all contracts/benefits allow coverage of healthcare professional administered drugs as part of their pharmacy benefit.
 - Not all contracts/benefits cover all medical infusible drugs.
- Unless otherwise indicated within drug specific criteria, the drugs listed in this policy are administered by a healthcare professional and therefore are covered under the medical benefit.
- Clinical documentation must be submitted for each request (initial and recertification [if applicable]) unless otherwise specified (e.g., provider attestation required). Supporting documentation includes, but is not limited to, progress notes documenting previous treatments and treatment history, diagnostic testing, laboratory test results, genetic testing or biomarker results, imaging, and other objective or subjective measures of clinical benefit.

For recertification, continued approval requires documentation demonstrating that the requested product is providing ongoing benefit to the patient, evidenced by improvement or stability in the disease state or condition, and that continued use remains medically necessary. Ongoing use of the requested product must continue to reflect the current policy's preferred formulary. Recertification reviews may result in the requirement to trial more cost-effective treatment alternatives as they become available (i.e., generics, biosimilars, or other guideline supported treatment options). Requested dosing must remain consistent with the FDA-approved or off-label/guideline-supported dosing recommendations.
- All requests will be reviewed to ensure they are being used for an appropriate indication and may be subject to an off-label review in accordance with our Off-Label Use of FDA Approved Drugs Policy (Pharmacy-32).
- Dose and frequency should be in accordance with the FDA label or recognized compendia (for off-label uses). When services are performed in excess of established parameters, they may be subject to review for medical necessity.
- All utilization management requirements outlined in this policy are compliant with applicable New York State insurance laws and regulations. Policies will be reviewed and updated as necessary to ensure ongoing compliance with all state and federally mandated coverage requirements.
- This policy does not apply to Medicare Part D and D-SNP pharmacy benefits. The drugs in this policy may apply to all other lines of business including Medicare Advantage.
- For members with Medicare Advantage, medications with a National Coverage Determination (NCD) and/or Local Coverage Determination (LCD) will be covered pursuant to the criteria outlined by the NCD and/or LCD. NCDs/LCDs for applicable medications can be found on the CMS website at <https://www.cms.gov/medicare-coverage-database/search.aspx>. In the absence of a Medicare National Coverage Determination (NCD), Local Coverage Determination (LCD), or other Medicare coverage guidance, CMS permits a Medicare Advantage Organization (MAO) to establish its own coverage determinations in accordance with 42 CFR § 422.101(b)(6). Indications that have not been addressed by the applicable medication's LCD/NCD will be covered in accordance with criteria

Pharmacy Management Drug Policy

Ophthalmic Vascular Endothelial Growth Factor (VEGF) Inhibitors

- determined by the Health Plan (which may include review per the Health Plan's Off-Label Use of FDA Approved Drugs policy). Step therapy requirements may be imposed in addition to LCD/NCD requirements.
9. Manufacturers may either discontinue participation in, or may not participate in, the Medicaid Drug Rebate Program (MDRP). Under New York State Medicaid requirements, physician-administered drugs must be produced by manufacturers that participate in the MDRP. Products made by manufacturers that do not participate in the MDRP will not be covered under Medicaid Managed Care/HARP lines of business. Drug coverage will not be available for any product from a non-participating manufacturer. For a complete list of New/Reinstated & Terminated Labelers please visit: <https://www.medicaid.gov/medicaid/prescriptiondrugs/medicaid-drug-rebate-program/newreinstated-terminated-labeler-information/index.html>
 10. The requested site of care may impact approval timeframe and subject to review.
 11. This policy is based on available evidence as of the last review date. Coverage determinations are subject to applicable plan documents, state and federal regulations, and individual patient circumstances. This policy does not constitute medical advice.
 12. For commercial contracts, medical necessity determinations align with the Certificate of Coverage issued by the Health Plan, which states that covered services must be clinically appropriate and not primarily for the convenience of the member, the member's family, or the provider.
 13. This policy is subject to ongoing revision. Newly marketed drugs and existing drugs with new indications may be subject to prior authorization until formal coverage criteria are established. Inclusion of a drug in this policy does not guarantee its current availability on the market, as some agents may be discontinued, withdrawn, or otherwise unavailable. As product status changes, drugs may be removed from the policy.

CODES:

Eligibility for reimbursement is based upon the benefits set forth in the member's subscriber contract. CODES MAY NOT BE COVERED UNDER ALL CIRCUMSTANCES. PLEASE READ THE POLICY AND GUIDELINES STATEMENTS CAREFULLY.

Codes may not be all inclusive as the AMA and CMS code updates may occur more frequently than policy updates.

Code Key:

Experimental/Investigational = (E/I),

Not medically necessary/ appropriate = (NMN).

Copyright © 2006 American Medical Association, Chicago, IL

HCPCS: Number Description

J0179	Beovu
Q5124	Byooviz
Q5128	Cimerli
J0178	Eylea
J0177	Eylea HD
J2778	Lucentis
Q5168	Nufymco
Q5147	Pavblu
J2779	Susvimo
J2777	Vabysmo
Q5155	Yesafili

Pharmacy Management Drug Policy

Ophthalmic Vascular Endothelial Growth Factor (VEGF) Inhibitors

UPDATES:

Date	Revision
08/07/2026	Revised
07/02/2026	Revised
06/01/2026	Revised
04/15/2026	Effective and Posted
11/13/2025	P&T Committee Approval

REFERENCES:

In addition to the full prescribing information for each individual drug, any references that have been utilized will be listed below.

1. Vemulakonda GA, Bailey ST, Kim SJ, et al. Age-Related Macular Degeneration Preferred Practice Pattern®. *Ophthalmology*. 2025;132(4):P1-P74. doi:10.1016/j.ophtha.2024.12.018
2. Wong WL, Su X, Li X, et al. Global prevalence of age-related macular degeneration and disease burden projection for 2020 and 2040: a systematic review and meta-analysis. *Lancet Glob Health*. 2014;2(2):e106-e116. doi:10.1016/S2214-109X(13)70145-1
3. Ferrara N, Adamis AP. Ten years of anti-vascular endothelial growth factor therapy. *Nat Rev Drug Discov*. 2016;15(6):385-403. doi:10.1038/nrd.2015.17
4. American Diabetes Association Professional Practice Committee. 12. Retinopathy, Neuropathy, and Foot Care: Standards of Care in Diabetes-2025. *Diabetes Care*. 2025;48(1 Suppl 1):S252-S265. doi:10.2337/dc25-S012
5. Diabetic Retinopathy Clinical Research Network, Wells JA, Glassman AR, et al. Aflibercept, bevacizumab, or ranibizumab for diabetic macular edema. *N Engl J Med*. 2015;372(13):1193-1203. doi:10.1056/NEJMoa1414264
6. Campochiaro PA, Heier JS, Feiner L, et al. Ranibizumab for macular edema following branch retinal vein occlusion: six-month primary end point results of a phase III study. *Ophthalmology*. 2010;117(6):1102-1112.e1. doi:10.1016/j.ophtha.2010.02.021
7. Tufail A, Narendran N, Patel PJ, et al. Ranibizumab in myopic choroidal neovascularization: the 12-month results from the REPAIR study. *Ophthalmology*. 2013;120(9):1944-5.e1. doi:10.1016/j.ophtha.2013.06.010
8. Mintz-Hittner HA, Kennedy KA, Chuang AZ; BEAT-ROP Cooperative Group. Efficacy of intravitreal bevacizumab for stage 3+ retinopathy of prematurity. *N Engl J Med*. 2011;364(7):603-615. doi:10.1056/NEJMoa1007374
9. Ekinci DY, Çelik K. Comparison of the Efficacy Between Intravitreal Aflibercept and Laser Photocoagulation in the Treatment of Retinopathy of Prematurity. *J Pediatr Ophthalmol Strabismus*. 2020;57(1):54-60. doi:10.3928/01913913-20191104-01
10. Stahl A, Lepore D, Fielder A, et al. Ranibizumab versus laser therapy for the treatment of very low birthweight infants with retinopathy of prematurity (RAINBOW): an open-label randomised controlled trial. *Lancet*. 2019;394(10208):1551-1559. doi:10.1016/S0140-6736(19)31344-3
11. Fogli S, Del Re M, Rofi E, Posarelli C, Figus M, Danesi R. Clinical pharmacology of intravitreal anti-VEGF drugs. *Eye (Lond)*. 2018;32(6):1010-1020. doi:10.1038/s41433-018-0021-7
12. Glassman AR. Anti-Vascular Endothelial Growth Factor Options and Questions for Diabetic Eye Disease Treatment. *JAMA Ophthalmol*. 2025;143(4):336-337. doi:10.1001/jamaophthalmol.2025.0007
13. Novartis Pharmaceuticals Corporation. *BEOVU® (brolucizumab-dbl) injection, for intravitreal use: Highlights of Prescribing Information*. East Hanover, NJ; Novartis: 2024. Accessed November 2025. Available from: https://www.novartis.com/us-en/sites/novartis_us/files/beovu.pdf
14. Genentech, Inc. *VABYSMO® (faricimab-svdn) injection for intravitreal use: Highlights of Prescribing Information*. South San Francisco, CA: Genentech, Inc.; 2024. Accessed November 2025. Available from: https://www.gene.com/download/pdf/vabysmo_prescribing.pdf
15. Bauman CR, Spaide RF, Vajzovic L, et al. Retinal Vasculitis and Intraocular Inflammation after Intravitreal Injection of Brolucizumab. *Ophthalmology*. 2020;127(10):1345-1359. doi:10.1016/j.ophtha.2020.04.017

Pharmacy Management Drug Policy

Ophthalmic Vascular Endothelial Growth Factor (VEGF) Inhibitors

16. Holekamp NM, Campochiaro PA, Chang MA, et al. Archway Randomized Phase 3 Trial of the Port Delivery System with Ranibizumab for Neovascular Age-Related Macular Degeneration. *Ophthalmology*. 2022;129(3):295-307. doi:10.1016/j.ophtha.2021.09.016
17. Jhaveri CD, Glassman AR, Ferris FL, et al. Aflibercept Monotherapy or Bevacizumab First for Diabetic Macular Edema. *N Engl J Med*. 2022;387(8):692-703. doi: 10.1056/NEJMoa2204225