

MEDICAL POLICY

MEDICAL POLICY DETAILS	
Medical Policy Title	Transcatheter Heart Valve Procedures
Policy Number	7.01.109
Category	Technology Assessment
Original Effective Date	07/15/22
Committee Approval Date	03/24/22, 03/23/23
Current Effective Date	03/23/23
Archived Date	N/A
Archive Review Date	N/A
Product Disclaimer	<ul style="list-style-type: none"> • If a product excludes coverage for a service, it is not covered, and medical policy criteria do not apply. • If a commercial product (including an Essential Plan or Child Health Plus product), medical policy criteria apply to the benefit. • If a Medicaid product covers a specific service, and there are no New York State Medicaid guidelines (eMedNY) criteria, medical policy criteria apply to the benefit. • If a Medicare product (including Medicare HMO-Dual Special Needs Program (DSNP) product) covers a specific service, and there is no national or local Medicare coverage decision for the service, medical policy criteria apply to the benefit. • If a Medicare HMO-Dual Special Needs Program (DSNP) product DOES NOT cover a specific service please refer to the Medicaid Product coverage line.

POLICY STATEMENT

I. AORTIC VALVE:

- A. Based upon our criteria and assessment of the peer-reviewed literature, transcatheter aortic valve implantation (TAVI) also known as transcatheter aortic valve replacement (TAVR) with an U.S. Food and Drug Administration (FDA) approved transcatheter heart valve system, performed via an approach consistent with the device's FDA-approved labeling, may be considered **medically necessary** for patients with native valve aortic stenosis and any surgical risk level when **ALL** of the following conditions are present:
1. Severe aortic stenosis (refer to Policy Guideline I) with a calcified aortic annulus; **AND**
 2. New York Heart Association heart failure class II, III, or IV symptoms; **AND**
 3. Left ventricular ejection fraction greater than 20%; **AND**
 4. Patient does not have a unicuspid aortic valves.
- B. Based upon our criteria and assessment of the peer-reviewed literature, transcatheter aortic valve implantation (TAVI) with a transcatheter heart valve system approved for use for repair of a degenerated bioprosthetic valve (valve-in-valve) may be considered **medically necessary** when **ALL** of the following conditions are present:
1. Failure (stenosed, insufficient, or combined) of a surgical bioprosthetic aortic valve; **AND**
 2. New York Heart Association heart failure class II, III, or IV symptoms; **AND**
 3. Left ventricular ejection fraction greater than 20%; **AND**
 4. Patient is not an operable candidate for open surgery, as judged by at least two cardiovascular specialists (cardiologist and/or cardiac surgeon); or patient is an operable candidate but is at high risk for open surgery.
- C. Based upon our criteria and assessment of the peer-reviewed literature, transcatheter aortic valve implantation (TAVI) has not been medically proven to be effective and, therefore, is considered **investigational** for all other indications.

II. MITRAL VALVE:

Medical Policy: Transcatheter Heart Valve Procedures

Policy Number: 7.01.109

Page 2 of 17

- A. Based upon our criteria and assessment of the peer-reviewed literature, transcatheter mitral valve repair (TMVR) with an U.S. Food and Drug Administration (FDA) approved device, may be considered **medically necessary** for patients with symptomatic, primary mitral regurgitation (MR), who are considered at prohibitive risk for open surgery (refer to Policy Guideline I).
- B. Based upon our criteria and assessment of the peer-reviewed literature, TMVR with an FDA approved device, may be considered **medically necessary** for patients with heart failure and moderate-to-severe or severe symptomatic secondary mitral regurgitation, despite the use of maximally tolerated guideline-directed medical therapy (refer to Policy Guideline IV).
- C. Based upon our criteria and assessment of the peer-reviewed literature, TMVR is considered **investigational** for all other indications.

III. PULMONARY VALVE:

- A. Based upon our criteria and assessment of the peer-reviewed literature, transcatheter pulmonary valve implantation (TPVI) with an U.S. Food and Drug Administration (FDA) approved valve, may be considered **medically necessary** for patients with congenital heart disease and current right ventricular outflow tract obstruction (RVOT) or regurgitation with one of the following indications:
 - 1. Individuals with right ventricle-to-pulmonary artery conduit with or without bioprosthetic valve with at least moderate pulmonic regurgitation;
 - 2. Individuals with native or patched RVOT with at least moderate pulmonic regurgitation;
 - 3. Individuals with right ventricle-to-pulmonary artery conduit with or without bioprosthetic valve with pulmonic stenosis (mean RVOT gradient at least 35 mm Hg); or
 - 4. Individuals with native or patched RVOT with pulmonic stenosis (mean RVOT gradient at least 35 mm Hg).
- B. Based upon our criteria and assessment of the peer-reviewed literature, TPVI is considered **investigational** for all other indications.

IV. TRICUSPID VALVE:

- A. Based upon our criteria and the lack of peer-reviewed literature, transcatheter tricuspid valve repair or replacement (TTVR) is considered **investigational** for all indications.
- V. Based upon our criteria and the lack of peer-reviewed literature, transcatheter placement and removal of a cerebral embolic protection device is considered **investigational**.

Refer to Corporate Medical Policy #11.01.03 Experimental and Investigational Services.

POLICY GUIDELINES

- I. The STS Short-Term Risk Calculator allows you to calculate a patient's risk of mortality and morbidities for the most commonly performed cardiac surgeries. Information regarding the risk calculator can be found at: <https://www.sts.org/resources/risk-calculator>.
- II. "Prohibitive Risk" for open surgery may be determined based on the presence of a Society for Thoracic Surgeons predicted mortality risk of 12% or greater .
- III. The U.S. Food and Drug Administration (FDA) Risk Levels for Open Surgery are defined as follows:
 - A. Extreme risk or inoperable for open surgery:
 - 1. Predicted risk of operative mortality and/or serious irreversible morbidity 50% or higher for open surgery.
 - B. High risk for open surgery:
 - 1. Society of Thoracic Surgeons predicted operative risk score of 8% or higher; or
 - 2. Judged by a heart team, which includes an experienced cardiac surgeon and a cardiologist, to have an expected mortality risk of 15% or higher for open surgery.
 - C. Intermediate risk for open surgery:
 - 1. Society of Thoracic Surgeons predicted operative risk score of 3% to 7%.
 - D. Low risk for open surgery:
 - 1. Society of Thoracic Surgeons predicted operative risk score of less than 3% or 4%.

Medical Policy: Transcatheter Heart Valve Procedures

Policy Number: 7.01.109

Page 3 of 17

IV. The New York Heart Association (NYHA) Heart Failure Classification (NYHA, 1994) are defined as follows:

NYHA Functional Class I: No limitation of physical activity. Ordinary physical activity does not cause undue fatigue, palpitation, dyspnea or anginal pain.

NYHA Functional Class II: Slight limitation of physical activity. Comfortable at rest. Ordinary physical activity results in fatigue, palpitation, dyspnea or anginal pain.

NYHA Functional Class III: Marked limitation of physical activity. Comfortable at rest. Less than ordinary activity causes fatigue, palpitation, dyspnea or anginal pain.

NYHA Functional Class IV: Unable to carry on any physical activity without discomfort. Symptoms of heart failure at rest. If any physical activity is undertaken, discomfort increases.

V. Optimal medical therapy include evaluation for cardiac risk factors (e.g., hypertension, diabetes mellitus, and hyperlipidemia), lifestyle factors (exercise, diet, smoking, and weight), risk for thromboembolic events, and drug therapy (e.g., vasodilators, diuretics, nitrates, nifedipine, and beta-adrenergic blocker therapy).

DESCRIPTION

Transcatheter Aortic Valve Implantation or Replacement

Aortic stenosis is defined as narrowing of the aortic valve opening, resulting in obstruction of blood flow from the left ventricle into the ascending aorta. Progressive calcification of the aortic valve is the most common etiology in North America and Europe, while rheumatic fever is the most common etiology in developing countries. Congenital abnormalities of the aortic valve (most commonly a bicuspid or unicuspid valve) increase the risk of aortic stenosis; however, aortic stenosis can also occur in a normal aortic valve. Risk factors for calcification of a congenitally normal valve mirror those for atherosclerotic vascular disease which include advanced age, male gender, smoking, hypertension, and hyperlipidemia. For symptomatic patients with severe aortic valve stenosis, the open-heart approach for surgical aortic valve replacement (SAVR) is currently the gold standard treatment. Long-term results are convincing, and even in octogenarians, SAVR is feasible with acceptable results. However, in patients with many co-morbidities, the outcome is less favorable, and many of these patients may be inoperable or carry an unacceptably high peri-operative risk.

Transcatheter aortic valve implantation (TAVI), also known as transcatheter aortic valve replacement (TAVR), represents an alternative to SAVR in patients who are elderly, inoperable or at high-risk for conventional surgery. TAVI is performed percutaneously, most often via the transfemoral artery approach but can also be done through the subclavian artery approach and transapically using mediastinoscopy. Balloon valvuloplasty is first performed to open up the stenotic area followed by passage of a bioprosthetic artificial valve across the native aortic valve. The valve is initially compressed to allow passage across the native valve and is then expanded and secured to the underlying aortic valve annulus. The procedure is performed on the beating heart without cardiopulmonary bypass. Once the prosthetic valve is deployed, angiography, computed tomography (CT) angiography or echocardiography is performed to ensure successful implantation of the device.

The currently available transcatheter aortic valves approved by the U.S. Food and Drug Administration (FDA) include the balloon-expandable Edwards Sapien 3 and Edwards Sapien XT (Edwards Lifesciences, Irvine, CA) and self-expandable Medtronic Evolt R and Evolut PRO systems (Medtronic, Inc., Santa Rosa, CA). They are indicated for percutaneous aortic valve implantation in individuals with severe aortic stenosis who are judged by a heart team, including a cardiac surgeon, to be high risk or inoperable for open aortic valve replacement. In September 2021, the FDA approved the Abbott Portico with FlexNav transcatheter aortic valve replacement (TAVR) system to treat people with symptomatic, severe aortic stenosis who are at high or extreme risk for open-heart surgery. There are several TAVI/transcatheter aortic valve replacement (TAVR) systems currently being studied which have not received FDA approval to include: Acurate TA transaortic valve replacement system; Engager TAVI system and JenaValve transapical (TAVI) system.

Medical Policy: Transcatheter Heart Valve Procedures

Policy Number: 7.01.109

Page 4 of 17

Transcatheter Mitral Valve Repair

Mitral Regurgitation (MR) is the second most common valvular heart disease, occurring in 7% of people, 75 years and older, and accounting for 24% of all patients with valvular heart disease. MR with accompanying valvular incompetence leads to left ventricular (LV) volume overload with secondary ventricular remodeling, myocardial dysfunction, and left heart failure. Clinical signs and symptoms of dyspnea and orthopnea may also be present in patients with valvular dysfunction. MR severity is classified as mild, moderate, or severe disease on the basis of echocardiographic and/or angiographic findings (1+, 2+, and 3 to 4+ angiographic grade, respectively).

Patients with MR generally fall into two categories, primary (also called degenerative) and secondary (also called functional) MR. Primary MR results from a structural abnormality in the valve, which causes it to leak. This leak may be a result of a floppy, or prolapsed leaflet, or a ruptured cord, which caused the leaflet to detach partially causing the leaflet to flail. Because the primary cause is a structural abnormality, most cases of primary MR are surgically corrected. Secondary MR results from LV dilatation due to ischemic or dilated cardiomyopathy. This causes the mitral valve (MV) leaflets not to coapt or meet in the center. Because the valves are structurally normal in secondary MR, correcting the dilated LV using medical therapy is the primary treatment strategy used in the United States.

Standard open MV repair requires thoracotomy and cardiopulmonary bypass, which may not be tolerated in the elderly or in patients with underlying cardiac disease or other co-morbid conditions. TMVR is a less invasive alternative to open surgical therapy. The MitraClip system consists of a catheter, a steerable sleeve, and the clip, which is 4-mm wide, and made of a cobalt-chromium alloy and polypropylene fabric. The MitraClip is deployed percutaneously via a transfemoral approach, with a transseptal puncture used to access the left side of the heart and the mitral valve. Placement of MitraClip leads to coaptation (joining) of the mitral leaflets, which creates a double-orifice valve.

In October 2013, the MitraClip Clip Delivery System (Abbott Vascular) was approved by the FDA through the premarket approval process for treatment of "significant symptomatic mitral regurgitation (MR 3+ or greater) due to primary abnormality of the mitral apparatus (degenerative MR) in patients who have been determined to be at a prohibitive risk for mitral valve surgery by a heart team. FDA product code: NKM.

In March 2019, the FDA approved a new indication for MitraClip, for "treatment of patients with normal mitral valves who develop heart failure symptoms and moderate-to-severe or severe mitral regurgitation because of diminished left heart function (commonly known as secondary or functional mitral regurgitation) despite being treated with optimal medical therapy. Optimal medical therapy includes combinations of different heart failure medications along with, in certain patients, cardiac resynchronization therapy and implantation of cardioverter defibrillators."

Transcatheter Pulmonary Valve Implantation

Congenital heart disease, including tetralogy of Fallot, pulmonary atresia, and transposition of the great arteries, is generally treated by surgical repair at an early age. This involves reconstruction of the right ventricular outflow tract (RVOT) and pulmonary valve using a surgical homograft or a bovine-derived valved conduit. These repairs are prone to the development of pulmonary stenosis or regurgitation over long periods of follow-up. Because individuals with surgically corrected congenital heart disease repair are living into adulthood, RVOT dysfunction following initial repair has become more common. Calcification of the RVOT conduit can lead to pulmonary stenosis, while aneurysmal dilatation can result in pulmonary regurgitation. RVOT dysfunction can lead to decreased exercise tolerance, potentially fatal arrhythmias, and/or irreversible right ventricular dysfunction.

Transcatheter pulmonary valve implantation (TPVI) is a less invasive alternative to open surgical pulmonary valve replacement or reconstruction for right ventricular outflow tract (RVOT) obstruction. Percutaneous pulmonary valve replacement may be indicated for congenital pulmonary stenosis. Pulmonary stenosis or regurgitation in a patient with congenital heart disease (CHD) who has previously undergone RVOT surgery are additional indications. Patients with prior CHD repair are at risk of needing repeated reconstruction procedures.

The Melody Transcatheter Pulmonary Valve (TPV) and the Ensemble Transcatheter Valve Delivery System are used together for percutaneous replacement of a dysfunctional pulmonary valve. The Melody valve consists of a section of bovine jugular vein with an intact native venous valve. The transcatheter delivery system consists of a balloon-in-balloon catheter with a retractable sheath and distal cup into which the valve is placed. The procedure is performed on a beating

Medical Policy: Transcatheter Heart Valve Procedures

Policy Number: 7.01.109

Page 5 of 17

heart without the use of cardiopulmonary bypass. In January 2010, the Melody TPV and the Ensemble Transcatheter Valve Delivery System (Medtronic) were approved by FDA under the HDE program and more recently in February of 2017 the FDA approval of the Melody system was expanded to include patients with a dysfunctional surgical bioprosthetic valve (valve-in-valve).

The Edwards SAPIEN XT Transcatheter Heart Valve (Pulmonic) (Edwards Lifesciences) is composed of a stainless-steel frame with bovine pericardial tissue leaflets and available in 23- and 26-mm sizes. It includes a delivery accessories system. On February 29, 2016, it was approved by FDA as a supplement "for use in pediatric and adult patients with a dysfunctional, noncompliant Right Ventricular Outflow Tract (RVOT) conduit with a clinical indication for intervention and either pulmonary regurgitation \geq moderate and/or mean RVOT gradient \geq 35 mmHg."

The Medtronic Harmony Transcatheter Pulmonary Valve (TPV) System is the first FDA-approved (March 2021) transcatheter valve system specifically designed for use in the management of pediatric and adult patients with severe pulmonary regurgitation who have a native or surgically repaired right ventricular outflow tract (RVOT) and are clinically indicated for surgical pulmonary valve replacement.

Transcatheter Tricuspid Valve Repair or Replacement

Clinically significant tricuspid regurgitation (TR) is quite common with at least moderate TR occurring in greater than one in 200 of the general population and four percent of those greater than or equal to 75 years of age. The underlying etiology of TR is most commonly pulmonary hypertension, either from left-sided heart failure, mitral or aortic valve disease, or primary pulmonary causes. Atrial fibrillation may be both a marker of disease progression as well as a cause of annular dilatation due to atrial remodeling. Neither medical therapy nor conventional surgery is efficacious for most patients with significant TR. Currently, tricuspid valve surgery for functional TR is recommended only when performing surgery for concomitant left-sided valve disease. Devices for transcatheter tricuspid valve repair (TTVR) or replacement are in early stages of development for the treatment of tricuspid regurgitation and there are clinical studies evaluating the use of TTVR devices. Individual selection criteria for percutaneous tricuspid valve replacement are based on limited data.

Currently, there are no FDA-approved devices for transcatheter tricuspid valve repair (TTVR) or replacement.

The 2017 position statement of the European Society of Cardiology Working Groups of Cardiovascular Surgery and Valvular Heart Disease: Management of Tricuspid Valve Regurgitation states that "Percutaneous tricuspid valve intervention (both repair and replacement) is still in its infancy but may become a reliable option in future, especially for high-risk patients with isolated primary TR or with secondary TR related to advanced left sided heart valve disease."

RATIONALE

Transcatheter heart valve replacement is a less invasive alternative to conventional open-heart surgery as it does not require heart-lung bypass. A catheter inserted using a transfemoral (TF), transapical or transaortic approach allows the introduction of an expandable prosthetic heart valve which is then delivered to the diseased native valve. The TF vascular access approach has been associated with reduced vascular complications (Carrol, 2020). The 2020 ACC/AHA guideline (Otto, 2020) recommendations for TAVR in moderate or lower STS risk patients specify that the TF vascular access approach should be used. Registry data shows that more than 90 percent of TAVR in the U.S. is now performed with the TF approach. Two minimally invasive alternatives to surgical mitral valve repair include transcatheter leaflet repair and percutaneous annuloplasty. The purpose of transcatheter mitral valve leaflet repair is to keep the two valve leaflets more closely fitted together, thereby reducing regurgitation. Percutaneous annuloplasty attempts to reshape the mitral annulus using catheters guided through the vasculature to reach the heart to reduce regurgitation.

Transcatheter Aortic Valve Implantation or Replacement

TAVR is well established for the treatment of high-risk and inoperable patients with symptomatic severe aortic stenosis. A robust evidence base has compared transcatheter aortic valve replacement (TAVR) to the standard of care for aortic stenosis. The series of Placement of AoRTic TraNscathetER Valves (PARTNER) trials began with PARTNER 1B (n=358), which demonstrated superiority of TAVR to medical therapy in inoperable patients, with an absolute survival advantage of 23percent at five years (Leon et al., 2010). The PARTNER 1A (n=699) and CoreValve (n=795) trials randomized high-surgical risk patients between TAVR and surgical aortic valve replacement (SAVR) (Smith et al., 2011;

Medical Policy: Transcatheter Heart Valve Procedures

Policy Number: 7.01.109

Page 6 of 17

Adams et al., 2014). Both trials were noninferiority trials and showed either no difference or improved survival with TAVR at one year. Patients in PARTNER 1A have been followed to five years with no survival difference seen (Sanchez et al., 2020; Reardon et al., 2019; Herrmann et al., 2019; Pibarot et al., 2019; Mack et al., 2015 and Kapadia et al., 2015).

Two multicenter randomized controlled studies have compared TAVR to surgical aortic valve replacement (SAVR) in symptomatic patients at intermediate surgical risk. The PARTNER 2A trial (Leon et al., 2016) randomized 2,032 patients to the balloon-expandable Sapien valve versus SAVR, and the SURTAVI trial (Reardon et al., 2017) randomized 1,660 patients to a self-expanding TAVR (CoreValve or Evolut-R) versus SAVR. The results of both trials demonstrated noninferiority of TAVR to SAVR for the composite endpoint of death and stroke at two years. In a large registry of symptomatic, intermediate-risk patients who underwent TAVR using the balloon-expandable Sapien 3 system (Thourani et al., 2016), survival was markedly superior to the surgical arm of the PARTNER 2A study (Herrmann et al., 2019).

Favorable short-term results with TAVR in low-risk patients were reported in two recent randomized clinical trials. The Evolut Low Risk Trial (Popma et al., 2019) reported the estimated two-year incidence of the primary endpoint, a composite of death or disabling stroke, was 5.3 percent in the TAVI group and 6.7 percent in the SAVR group showing non-inferiority of TAVI and SAVR, but no superiority for either mortality or stroke at one year. The PARTNER 3 Trial (Mack et al., 2019) low-risk patient study showed superiority of TAVI for stroke and the composite primary endpoint of death, stroke and rehospitalization at one year. The Nordic Aortic Valve Intervention Trial (NOTION) (Thyregod et al., 2019) randomized patients to receive TAVR or SAVR, and 82 percent of the patients were at low risk for surgical operations (i.e., Society of Thoracic Surgeons Predicted Risk of Mortality [STS-PROM] score less than 4 percent).

Similar outcomes were achieved in both TAVR and SAVR treatment arms at five years. Wakesman et al. (2018) reported in a prospective study that transfemoral TAVR, using mainly a third-generation balloon-expandable TAVR device, was associated with no deaths at 30 days compared with 1.7 percent in a historical, propensity-matched SAVR cohort. The risk/benefit profile for periprocedural complications in low-risk patients is similar to the overall TAVR population (i.e., reduction in acute kidney injury and bleeding on the one hand and an increase in pacemaker implantation and vascular complications) (Overtchouk et al., 2019). The 2017 focused update to the 2014 American College of Cardiology (ACC) and American Heart Association (AHA) Practice Guideline for the Management of Patients with Valvular Heart Disease recommends surgical aortic valve replacement for low surgical risk patients (Nishimura et al., 2017). However, long-term follow-up data on outcomes and valve durability is needed.

In June 2019, the National Institute for Health and Care Excellence published interventional procedure guidance regarding valve-in-valve TAVI for aortic bioprosthetic valve dysfunction. The guidance recommendation is that "Current evidence on the safety and efficacy of valve-in-valve transcatheter aortic valve implantation for aortic bioprosthetic dysfunction is adequate to support the use of this procedure provided that standard arrangements are in place for clinical governance, consent and audit."

In 2020, a new full ACC/AHA guideline was published that replaces the 2014 revision and 2017 focused update. The 2020 guidelines made recommendations on timing of intervention and choice of surgical or transcatheter intervention for treatment of aortic stenosis. Additionally, the guidelines state the following:

- "Treatment of severe aortic stenosis with either a transcatheter or surgical valve prosthesis should be based primarily on symptoms or reduced ventricular systolic function. Earlier intervention may be considered if indicated by results of exercise testing, biomarkers, rapid progression, or the presence of very severe stenosis."
- "Indications for TAVI are expanding as a result of multiple randomized trials of TAVI versus surgical aortic valve replacement. The choice of type of intervention for a patient with severe aortic stenosis should be a shared decision-making process that considers the lifetime risks and benefits associated with type of valve (mechanical versus bioprosthetic) and type of approach (transcatheter versus surgical)."

Transcatheter Mitral Valve Repair

Feldman et al. (2015) reported on the results of EVEREST II, an RCT that evaluated symptomatic or asymptomatic patients with grade 3+ or 4+ chronic MR who had SMR or primary MR etiology to TMVR at five years; patients were randomized to MitraClip or open MV repair/replacement. Most patients (73%) had primary MR and were eligible for

Medical Policy: Transcatheter Heart Valve Procedures

Policy Number: 7.01.109

Page 7 of 17

mitral repair or replacement surgery. Results showed that TMVR was less effective at reducing MR and of subsequent

Medical Policy: Transcatheter Heart Valve Procedures

Policy Number: 7.01.109

Page 8 of 17

surgery for MV dysfunction than conventional surgery, few patients experienced worsening MR or surgery after 6-month follow-up. There was no difference in long-term survival after TMVR compared to surgery and no decrement in left ventricle (LV) systolic function. Long-term survival was similar between the two treatment arms; however, functional MR and advanced age were associated with decreased survival regardless of the treatment.

The evidence for the use of MitraClip in patients with secondary mitral regurgitation (SMR) consists of two randomized-controlled-trials (RCTs), the Cardiovascular Outcomes Assessment of the MitraClip Percutaneous Therapy for Heart Failure Patients with Functional Mitral Regurgitation (COAPT), and the Percutaneous Repair with the MitraClip Device for Severe Functional/Secondary Mitral Regurgitation (MITRA-FR). Both trials compared MitraClip plus medical therapy to medical therapy alone in patients with SMR and heart failure, but they differed in their eligibility criteria, and primary outcome measures. COAPT enrolled 614 patients at 78 centers in the U.S. and Canada. MITRA-FR enrolled 304 patients at 37 centers in France. COAPT found a significant benefit for Mitraclip on the primary efficacy outcome (all HF hospitalizations within 24 months) and the primary safety outcome (freedom from device-related complications at 12 months). In contrast, the MITRA-FR investigators found no significant differences between Mitra-Clip plus medical therapy and medical therapy alone on the composite primary outcome (death from any cause or unplanned HF hospitalization at 12 months) or any secondary outcome, including all-cause mortality at 12 and 24 months and cardiovascular death at 12 and 24 months. Although the reasons for these discrepant results are not entirely clear, differences in the studies' design and conduct have been proposed as possible explanations. The severity of MR and heart failure among the patients in the trials differed. COAPT participants had more severe MR at baseline (effective regurgitant orifice area 41 versus 31 mm²) and remained symptomatic despite the use of maximal doses of guideline-directed medical therapy. In both trials, eligible patients had to be symptomatic despite the use of optimal medical therapy. In COAPT, however, a central eligibility committee confirmed that the patient was using maximal doses of guideline-directed medical therapy prior to enrollment, and patients who improved with medical therapy were excluded. MITRA-FR had less stringent eligibility criteria and patients had more changes in medical therapy during the trial, indicating their treatment might not have been optimized. Additionally, patients in MITRA-FR had further progressed heart failure as indicated by LV dilation and may have been less likely to benefit from MR treatment. There is some evidence that technical success and procedural safety differed between the trials. Procedural complications were higher in MITRA-FR than in COAPT, and more patients in MITRA-FR experienced residual MR class greater than 3+ post-procedure (both acutely and at 12 months).

A Report of the American College of Cardiology (ACC)/American Heart Association (AHA) Joint Committee on Clinical Practice Guidelines for the Management of Patients with Valvular Heart Disease (2020) recommendations for intervention for chronic primary MR regarding transcatheter edge-to-edge repair (TEER) include the following: (1) In severely symptomatic patients (NYHA class III or IV) with primary severe MR and high or prohibitive surgical risk, transcatheter edge-to-edge repair (TEER) is reasonable if mitral valve anatomy is favorable for the repair procedure

and patient life expectancy is at least one year: Class: 2a; Level of Evidence: B-R; (2) In symptomatic patients with severe primary MR attributable to rheumatic valve disease, mitral valve repair may be considered at a Comprehensive Valve Center by an experienced team when surgical treatment is indicated, if a durable and successful repair is likely: Class: 2b; Level of Evidence: B-R; (3) In patients with chronic severe secondary MR related to LV systolic dysfunction (LVEF less than 50%) who have persistent severe symptoms (NYHA class II, III, or IV) while on optimal guideline directed medical therapy (GDMT) for HF (Stage D), TEER is reasonable in patients with appropriate anatomy as defined on transesophageal echocardiography (TEE) and with LVEF between 20% and 50%, LVESD 70 mm or less, and pulmonary artery systolic pressure 70 mm Hg or less (Class: 2a; Level of Evidence: B-R). In summary, a mitral transcatheter edge-to-edge repair is of benefit to patients with severely symptomatic primary mitral regurgitation who are at high or prohibitive risk for surgery, as well as to a select subset of patients with secondary mitral regurgitation who remain severely symptomatic despite GDMT for heart failure. ACC/AHA guidelines for the management of patients with valvular heart disease (Nishimura et al., 2014) make the following recommendations regarding transcatheter valve replacement: Transcatheter mitral valve repair may be considered for severely symptomatic patients (NYHA class III to IV) with chronic severe primary MR (stage D) who have favorable anatomy for the repair procedure and a reasonable life expectancy but who have a prohibitive surgical risk because of severe comorbidities and remain severely symptomatic despite optimal guideline-directed medical therapy for heart failure. (Class IIb recommendation, level of evidence B -

Medical Policy: Transcatheter Heart Valve Procedures

Policy Number: 7.01.109

Page 9 of 17

procedure may be considered but usefulness/efficacy is less well established based on conflicting evidence from a single randomized trial or nonrandomized studies.)

Transcatheter Pulmonary Valve Implantation

Transcatheter pulmonary valve (TPV) placement was first reported in 2000. Beginning in January 2007, the Melody TPV (Medtronic, Inc., Santa Ana, CA) was implanted in 150 patients at five US centers under an Investigational Device Exemption (IDE) protocol for treatment of right ventricular outflow tract (RVOT) dysfunction. In January 2010, enrollment in the US Melody Valve IDE trial was completed, and the Melody valve was approved for placement in dysfunctional RVOT conduits as a palliative measure aimed at delaying surgical intervention (McElhinney et al., 2011). The trial was initially designed to follow patients for five years after implantation or until explantation but was modified in 2011 to allow follow-up out to 10 years in patients who provided supplemental written informed consent (Cheatham et al., 2015).

In January 2015, the Melody TPV received Pre-Market Approval (PMA) from the U.S. Food and Drug Administration (FDA) approval based on clinical evidence from three clinical studies that followed patients implanted with Melody TPV (i.e., the Melody U.S. IDE Study, the Melody U.S. Post Approval Study [PAS] and the European and Canadian Post-Market Surveillance Study [PMSS]). On February 24, 2017, approval of the Melody system was expanded to include patients with a dysfunctional surgical bioprosthetic valve (valve-in-valve). Per the FDA Summary of Effectiveness and Safety Data (SSED), the clinical data supporting the PMA supplemental approval decision were pooled from the following three (3) sources: Melody Transcatheter Pulmonary Valve (TPV) Long-term Follow-up Post Approval Study (PAS) n=8 patients; Melody TPV New Enrollment PAS n=17 patients and Real-World Data n=100 patients.

Cheatham et al. (2015) evaluated the midterm hemodynamic and clinical outcomes in the U.S. Melody Valve IDE trial patients (n=148), who were all at least four years out from Melody valve implantation. The nonrandomized IDE trial prospectively enrolled pediatric and adult patients (median age, 19 years) with right ventricular outflow tract conduit obstruction or regurgitation. The patients received and were discharged with a TPV were followed up annually according to a standardized protocol. During a median follow-up of 4.5 years (range, 0.4-7 years), 32 patients underwent right ventricular outflow tract reintervention for obstruction (n=27, with stent fracture in 22), endocarditis (n=3, 2 with stenosis and 1 with pulmonary regurgitation), or right ventricular dysfunction (n=2). Eleven patients had the TPV explanted as an initial or second reintervention. Five-year freedom from reintervention and explantation was 76±4 percent and 92±3 percent, respectively. A conduit present and lower discharge right ventricular outflow tract gradient were associated with longer freedom from reintervention. In the 113 patients who were alive and reintervention free, the follow-up gradient (median, 4.5 years after implantation) was unchanged from early post-TPV replacement, and all but one patient had mild or less pulmonary regurgitation. Almost all patients were in New York Heart Association class I or II. More severely impaired baseline spirometry was associated with a lower likelihood of improvement in exercise function after TPV replacement. The authors reported that TPV replacement with the Melody valve provided good hemodynamic and clinical outcomes up to seven years after implantation. Primary valve failure was rare. The main cause of TPV dysfunction was stenosis related to stent fracture, which was uncommon once presenting became more widely adopted.

In February 2016, the SAPIEN XT Transcatheter Heart Valve received Pre-Market Approval (PMA) from the U.S. Food and Drug Administration (FDA) approval based on clinical evidence from the Congenital Multicenter trial of Pulmonic Valve regurgitation Studying the SAPIEN IntervENTIONal (COMPASSION) THV trial. The 2016 FDA PMA approval states that Edwards agreed to conduct a study to evaluate long-term safety and effectiveness of the SAPIEN XT THV in the pulmonic position for the intended patient population (especially pediatric) when used as indicated with all valve sizes. It is a single-arm, prospective, multicenter post approval study using a performance goal based on the original COMPASSION trial. The study patients are pediatric and adult patients with a dysfunctional, non-compliant right ventricular outflow tract (RVOT) conduit with a clinical indication for intervention and pulmonary regurgitation greater than or equal to moderate and/or mean RVOT gradient greater than or equal to 35 mmHg. The eligibility criteria were consistent with the final FDA-approved IFU and labeling. A sample size of 162 subjects was required for the hypothesis test on the primary effectiveness endpoint with at least 80 percent of the power. A total of 191 patients were enrolled at up to 10 sites in the United States to account for loss to follow-up. The patients are being followed at hospital discharge: 30 days, one year and annually thereafter through five years.

Medical Policy: Transcatheter Heart Valve Procedures

Policy Number: 7.01.109

Page 10 of 17

The AHA/ACC Practice Guideline for the Management of Patients with Valvular Heart Disease does not make specific recommendations regarding the treatment of primary pulmonary valve disease (Nishimura et al, 2014, 2017).

The ACC/AHA 2008 Guidelines for the Management of Adults with Congenital Heart Disease does not include recommendations for transcatheter pulmonary valve implantation (Warnes et al., 2008). This guideline was updated in 2018 (Stout et al., 2018). The updated guideline addresses percutaneous pulmonary replacement with recommendations for Tetralogy of Fallot (TOF) stating:

- Pulmonary valve replacement (surgical or percutaneous) for relief of symptoms is recommended for patients with repaired TOF and moderate or greater pulmonary regurgitation (PR) with cardiovascular symptoms not otherwise explained (Class or Recommendation I; Level of evidence B-NR)
- Pulmonary valve replacement (surgical or percutaneous) is reasonable for preservation of ventricular size and function in asymptomatic patients with repaired TOF and ventricular enlargement or dysfunction and moderate or greater PR (Class or Recommendation IIa; Level of evidence B-NR)

Transcatheter Tricuspid Valve Repair or Replacement

Tricuspid valve repair or replacement via transcatheter approach devices are in early stages of development for the treatment of tricuspid regurgitation. There are small case series as well as ongoing clinical trials for patients with diseased tricuspid valves undergoing transcatheter tricuspid valve replacement. There is currently insufficient published evidence to assess the safety and/or impact on health outcomes of transcatheter tricuspid valve replacement in patients with diseased tricuspid valves.

Nickenig et al., (2019) report the 6-month safety and performance of a transcatheter tricuspid valve reconstruction system in the treatment of moderate to severe functional tricuspid regurgitation (TR) in 30 patients enrolled in the TRIREPAIR (TrIcuspid Regurgitation RePAIr with CaRdioband Transcatheter System) study. Between October 2016 and July 2017, 30 patients were enrolled in this single-arm, multicenter, prospective trial. Patients were diagnosed with moderate to severe, symptomatic TR in the absence of untreated left-heart disease and deemed inoperable because of unacceptable risk for open-heart surgery by the local heart team. Clinical, functional, and echocardiographic data were prospectively collected before and up to six months post-procedure. An independent core lab assessed all echocardiographic data, and an independent clinical event committee adjudicated the safety events. Mean patient age was 75 years, 73 percent were female, and 23 percent had ischemic heart disease. At baseline, 83 percent were in New York Heart Association (NYHA) functional class III to IV and mean left ventricular ejection fraction was 58 percent. Technical success was 100 percent.

Three patients died post procedure through six months. Between six months and baseline, echocardiography showed average reductions of annular septolateral diameter of 9 percent (42 mm vs. 38 mm; $p < 0.01$) proximal isovelocity surface area effective regurgitant orifice area of 50 percent (0.8 cm² vs. 0.4 cm²; $p < 0.01$) and mean vena contracta width of 28 percent (1.2 cm vs. 0.9 cm; $p < 0.01$). Clinical assessment showed that 76 percent of patients improved by at least 1 NYHA functional class with 88 percent in NYHA functional class I or II. Six-minute walk distance improved by 60 m ($p < 0.01$), and Kansas City Cardiomyopathy Questionnaire score improved by 24 points ($p < 0.01$). In conclusion, six-month outcomes show that the system performs as intended and appears to be safe in patients with symptomatic and moderate to severe functional TR. Significant reduction of TR through decrease of annular dimensions, improvements in heart failure symptoms, quality of life, and exercise capacity were observed. Further studies are warranted to validate these initial promising results.

Taramasso et al., (2019) reported on a large, prospective international registry which was developed to evaluate the initial clinical applications of transcatheter tricuspid valve intervention (TTVI) with different devices. TTVI for native tricuspid valve dysfunction has been emerging during the last few years as an alternative therapeutic option to serve a large high-risk population of patients with severe symptomatic tricuspid regurgitation (TR). The TriValve Registry included 312 high-risk patients with severe TR (76.4 +/- 8.5 years of age; 57% female; EuroSCORE II 9 +/- 8%) at 18 centers.

Interventions included repair at the level of the leaflets (MitraClip, Abbott Vascular, Santa Clara, California; PASCAL Edwards Lifesciences, Irvine, California), annulus (Cardioband, Edwards Lifesciences; TriCinch, 4tech, Galway, Ireland; Trialign, Mitraling, Tewksbury, Massachusetts), or coaptation (FORMA, Edwards Lifesciences) and replacement (Caval Implants, NaviGate, NaviGate Cardiac Structures, Lake Forest, California). Clinical outcomes were prospectively determined during mid-term follow-up. A total of 108 patients (34.6%) had prior left heart valve intervention (84 surgical

Medical Policy: Transcatheter Heart Valve Procedures

Policy Number: 7.01.109

Page 11 of 17

and 24 transcatheter, respectively). TR etiology was functional in 93 percent, and mean annular diameter was 46.9 +/- 9 mm. In 75 percent of the patients, the regurgitant jet was central (vena contracta 1.1 +/- 0.5; effective regurgitant orifice area 0.78 +/- 0.6 cm(2)). Pre-procedural systolic pulmonary artery pressure was 41 +/- 14.8 mm Hg. Implanted devices included: MitraClip in 210 cases, Trialign in 18 cases, TriCinch first generation in 14 cases, caval valve implantation in 30 cases, FORMA in 24 cases, Cardioband in 13 cases, NaviGate in six cases, and PASCAL in one case. In 64 percent of the cases, TTVI was performed as a stand-alone procedure. Procedural success was defined as successful device implantation and residual TR. The report concluded that TTVI is feasible with different technologies, has a reasonable overall procedural success rate, and is associated with low mortality and significant clinical improvement. Mid-term survival of this high-risk population is favorable. The report also noted that greater coaptation depth is associated with reduced procedural success, which is an independent predictor of mortality.

CODES

- Eligibility for reimbursement is based upon the benefits set forth in the member's subscriber contract.
- **CODES MAY NOT BE COVERED UNDER ALL CIRCUMSTANCES. PLEASE READ THE POLICY AND GUIDELINES STATEMENTS CAREFULLY.**
- Codes may not be all inclusive as the AMA and CMS code updates may occur more frequently than policy updates.
- Code Key: Experimental/Investigational = (E/I), Not medically necessary/ appropriate = (NMN).

CPT Codes

Code	Description
33361	Transcatheter aortic valve replacement (TAVR/TAVI) with prosthetic valve; percutaneous femoral artery approach
33362	open femoral artery approach
33363	open axillary artery approach
33364	open iliac artery approach
33365	transaortic approach (eg, median sternotomy, mediastinotomy)
33366	transapical exposure (eg, left thoracotomy)
33367	Transcatheter aortic valve replacement (TAVR/TAVI) with prosthetic valve; cardiopulmonary bypass support with percutaneous peripheral arterial and venous cannulation (eg, femoral vessels) (List separately in addition to code for primary procedure)
33368	Transcatheter aortic valve replacement (TAVR/TAVI) with prosthetic valve; cardiopulmonary bypass support with open peripheral arterial and venous cannulation (eg, femoral, iliac, axillary vessels) (List separately in addition to code for primary procedure)
33369	Transcatheter aortic valve replacement (TAVR/TAVI) with prosthetic valve; cardiopulmonary bypass support with central arterial and venous cannulation (eg, aorta, right atrium, pulmonary artery) (List separately in addition to code for primary procedure)
33370 (E/I)	Transcatheter placement and removal of a cerebral embolic protection device(s), including atrial access, catheterization, imaging, and radiological supervision and interpretation, percutaneous (List separately in addition to code for primary procedure)
33418	Transcatheter mitral valve repair, percutaneous approach, including transseptal puncture when performed; initial prosthesis
33419	Transcatheter mitral valve repair, percutaneous approach, including transseptal puncture when performed; additional prosthesis(es) during same session (List separately in addition to code for primary procedure)

Medical Policy: Transcatheter Heart Valve Procedures

Policy Number: 7.01.109

Page 12 of 17

33477	Transcatheter pulmonary valve implantation, percutaneous approach, including pre-stenting of the valve delivery site, when performed
0345T	Transcatheter mitral valve repair percutaneous approach via the coronary sinus
0483T	Transcatheter mitral valve implantation/replacement (TMVI) with prosthetic valve; percutaneous approach, including transseptal puncture, when performed
0544T	Transcatheter mitral valve annulus reconstruction, with implantation of adjustable annulus reconstruction device, percutaneous approach including transseptal puncture
0545T (E/I)	Transcatheter tricuspid valve annulus reconstruction with implantation of adjustable annulus reconstruction device, percutaneous approach
0569T (E/I)	Transcatheter tricuspid valve repair, percutaneous approach; initial prosthesis
0570T (E/I)	Transcatheter tricuspid valve repair, percutaneous approach; each additional prosthesis during same session (List separately in addition to code for primary procedure)
0646T (E/I)	Transcatheter tricuspid valve implantation (TTVI)/replacement with prosthetic valve, percutaneous approach, including right heart catheterization, temporary pacemaker insertion, and selective right ventricular or right atrial angiography, when performed

Copyright © 2023

American Medical Association, Chicago, IL

HCPCS Codes

Code	Description
No Codes	

ICD10 Codes

Code	Description
Multiple Codes	

REFERENCES

Aortic Valve:

*Adams DH, et al. U.S. CoreValve Clinical Investigators. Transcatheter aortic-valve replacement with a self-expanding prosthesis. N Engl J Med 2014 May 8;370(19):1790-8.

*Herrmann HC and Mack MJ. Transcatheter therapies for valvular heart disease. Bonow: Braunwald's heart disease-a textbook of cardiovascular medicine, 11th ed. Saunders, an imprint of Elsevier, 2019. Ch 72.1464-82.

Husso A, et al. Transcatheter and surgical aortic valve replacement in patients with bicuspid aortic valve. Clinical Research in Cardiology 2021;110:429-439.

*Kapadia SR, et al. PARTNER trial investigators. 5-year outcomes of transcatheter aortic valve replacement compared with standard treatment for patients with inoperable aortic stenosis (PARTNER 1): a randomized controlled trial. Lancet 2015 Jun 20;385(9986):2485-91.

Leon MB, et al. Outcomes two years after transcatheter aortic valve replacement in patients at low surgical risk. Journal of the American College of Cardiology 2021;77(9):1149-1161.

Medical Policy: Transcatheter Heart Valve Procedures

Policy Number: 7.01.109

Page 13 of 17

*Leon MB, et al. PARTNER Trial Investigators. Transcatheter aortic-valve implantation for aortic stenosis in patients who cannot undergo surgery. N Engl J Med 2010 Oct 21;363(17):1597-607.

*Leon MB, et al. PARTNER 2 Investigators. Transcatheter or surgical aortic-valve replacement in intermediate-risk patients. N Engl J Med 2016 Apr 28;374(17):1609-20.

*Mack MJ, et al. PARTNER 1 Trial Investigators. 5-year outcomes of transcatheter aortic valve replacement or surgical aortic valve replacement for high surgical risk patients with aortic stenosis (PARTNER 1): a randomised controlled trial. Lancet 2015 Jun 20;385(9986):2477-84.

Mack MJ, et al. PARTNER 3 Investigators. Transcatheter aortic-valve replacement with a balloon-expandable valve in low-risk patients. N Engl J Med 2019 May 2;380(18):1695-1705.

Makkar RR, et al. Five-year outcomes of transcatheter or surgical aortic-valve replacement. N Engl J Med 2020;382:799-809.

Michel JM, et al. Safety and efficacy of minimalist transcatheter aortic valve implantation using a new-generation balloon-expandable transcatheter heart valve in bicuspid and tricuspid aortic valves. Clin Res Cardiol Dec 2021;110(12):1993-2006. (Online Ahead of Print)

Mentias A, et al. Incidence, predictors, and outcomes of endocarditis after transcatheter aortic valve replacement in the United States. J Am Coll Cardiol Intv 2020;13(17):1973-1982.

National Institute for Health and Care Excellence. Interventional procedure overview of valve-in-valve TAVI for aortic bioprosthetic valve dysfunction (IP 1013/2 [IPG653]). June 2019.

[<https://www.nice.org.uk/guidance/ipg653/evidence/overview-final-pdf-6834685357>] Accessed 02/21/23.

National Institute for Health and Care Excellence. Valve-in-valve TAVI for aortic bioprosthetic valve dysfunction, Interventional procedures guidance [IPG653]. June 2019. [<https://www.nice.org.uk/guidance/ipg653>] Accessed 02/21/23.

*Nishimura RA, et al. 2017 AHA/ACC Focused update of the 2014 AHA/ACC guideline for the management of patients with valvular heart disease: a report of the American College of Cardiology/American Heart Association Task Force on clinical practice guidelines. J Am Coll Cardiol 2017 Jul 11;70(2):252-289.

*Nishimura RA, et al. 3rd. 2019 AATS/ACC/ASE/SCAI/STS expert consensus systems of care document: a proposal to optimize care for patients with valvular heart disease: a joint report of the American Association for Thoracic Surgery, American College of Cardiology, American Society of Echocardiography, Society for Cardiovascular Angiography and Interventions, and Society of Thoracic Surgeons. J Am Coll Cardiol 2019 May 28;73(20):2609-5.

*Otto CM, et al. 2020 ACC/AHA Guideline for the management of patients with valvular heart disease. Journal of the American College of Cardiology 2021;77(4).

[https://www.jacc.org/doi/pdf/10.1016/j.jacc.2020.11.018?_ga=2.237443847.627642434.1617300438-205757524.1586972163] Accessed 02/21/23.

*Overtchouk P, et al. Why should we extend transcatheter aortic valve implantation to low-risk patients? A comprehensive review. Arch Cardiovasc Dis 2019 May;112(5):354-362.

*Pibarot P and O’Gara PT. Prosthetic Heart Valves. Bonow: Braunwald’s heart disease-a textbook of cardiovascular medicine, 11th ed. Saunders, an imprint of Elsevier, 2019. Ch 71.1455-62.

*Popma JJ, et al. Evolut Low Risk Trial Investigators. Transcatheter aortic-valve replacement with a self-expanding valve in low-risk patients. N Engl J Med 2019 May 2;380(18):1706-1715.

*Reardon MJ, et al. SURTAVI Investigators. Surgical or transcatheter aortic-valve replacement in intermediate-risk patients. N Engl J Med 2017 Apr 6;376(14):1321-1331.

*Reardon MJ, et al. Reprise III Investigators. Two-year outcomes after transcatheter aortic valve replacement with mechanical versus self-expanding valves: The REPRISSE III randomized clinical trial. JAMA Cardiol 2019 Mar 1;4(3):223-229.

Medical Policy: Transcatheter Heart Valve Procedures

Policy Number: 7.01.109

Page 14 of 17

*Sanchez C, et al. Valvular heart disease. Conn's Current Therapy/Elsevier 2020:160-166.

*Smith CR, et al.; PARTNER Trial Investigators. Transcatheter versus surgical aortic-valve replacement in high-risk patients. N Engl J Med 2011 Jun 9;364(23):2187-98.

*Thourani VH, et al. Transcatheter aortic valve replacement versus surgical valve replacement in intermediate-risk patients: a propensity score analysis. Lancet 2016 May 28;387(10034):2218-25.

*Thyregod HGH, et al. Five-year clinical and echocardiographic outcomes from the nordic aortic valve intervention (NOTION) randomized clinical trial in lower surgical risk patients. Circulation 2019 Feb 1.

*Waksman R, et al. Transcatheter aortic valve replacement in low-risk patients with symptomatic severe aortic stenosis. J Am Coll Cardiol 2018 Oct 30;72(18):2095-2105.

Mitral Valve:

Ailawadi G, et al. One-year outcomes after MitraClip for functional mitral regurgitation. Circulation 2019 Jan 2;139(1):37-47.

Arnold SV, et al. Health status after transcatheter mitral-valve repair in heart failure and secondary mitral regurgitation; COAPT trial. J Am Coll Cardiol 2019 May 7;73(17):2123-2132.

Bonow RO, et al. 2019 AATS/ACC/SCAI/STS expert consensus systems of care document: operator and institutional recommendations and requirements for transcatheter mitral valve intervention: a joint report of the American Association for Thoracic Surgery, the American College of Cardiology, the Society for Cardiovascular Angiography and Interventions, and the Society of Thoracic Surgeons endorsed by the Heart Failure Society of America. J Thorac Cardiovasc Surg 2020 Jul;160(1):72-92.

Cubero-Gallego H, et al. Outcomes with percutaneous mitral repair vs optimal medical treatment for functional mitral regurgitation: a systematic review. Ann Transl Med 2020 Aug;8(15):962.

*Feldman T, et al. Randomized Comparison of Percutaneous Repair and Surgery for Mitral Regurgitation: 5-Year Results of EVEREST II. J Am Coll Cardiol 2015; 66(25): 2844-2854.

Kampaktsis PN, et al. Mitral regurgitation in 202: the 2020 focused update of the 2017 American College of Cardiology Expert Consensus Decision Pathway on the Management of Mitral Regurgitation. J Cardiothorac Vasc Anesth 2021 Jun;35(6):1678-1690.

Mack MJ, et al. 3-year outcomes of the transcatheter mitral valve repair in patients with heart failure. J Am Coll Cardiol 2021;77(8):1029-1040.

Obadia JF, et al. Percutaneous repair or medical treatment for secondary mitral regurgitation. N Engl J Med 2018;379(24):2297-2306.

Otto CM, et al. 2020 ACC/AHA Guideline for the management of patients with valvular heart disease: a report of the American College of Cardiology/American Heart Association Joint Committee on clinical practice guidelines. Circulation 2020;143(5):e35-71.

Otto CM, et al. 2020 ACC/AHA Guidelines for the management of patients with valvular heart disease: executive summary: a report of the American College of Cardiology/American Heart Association Joint Committee on clinical practice guidelines. J Am Coll Cardiol 2021;77(4):450-500.

Roques F, et al. The logistic EuroSCORE. Eur Heart J 2003 May;24(9):881-2.

Society for Thoracic Surgeons STS Short-Term Risk Calculator version 4.20. 2018 Nov 15.
[\[https://riskcalc.sts.org/stswebriskcalc/calculate\]](https://riskcalc.sts.org/stswebriskcalc/calculate) Accessed 02/21/23.

Stone GW, et al. Transcatheter mitral-valve repair in patients with heart failure. N Engl J Med 2018;379(24):2307-2318.

Pulmonary Valve:

Armstrong AK, et al. Association between patient age at implant and outcomes after transcatheter pulmonary valve replacement in the multicenter Melody valve trials. Catheter Cardiovasc Interv 2019 Oct 1;94(4):607-617.

Medical Policy: Transcatheter Heart Valve Procedures

Policy Number: 7.01.109

Page 15 of 17

Balzer D. Pulmonary valve replacement for tetralogy of fallot. Methodist Debaque Cardiovasc J 2019 Apr-Jun;15(2):122-132.

Cabalka AK, et al. Transcatheter pulmonary valve replacement using the melody valve for treatment of dysfunctional surgical bioprostheses: a multicenter study. J Thorac Cardiovasc Surg 2018 Apr;155(4):1712-1724.e1.

*Cheatham JP, et al. Clinical and hemodynamic outcomes up to 7 years after transcatheter pulmonary valve replacement in the US Melody Valve Investigational Device Exemption Trial. Circulation 2015 Jun 2;131(22):1960-70.

Hascoet S, et al. Early outcomes of percutaneous pulmonary valve implantation using the Edwards SAPIEN 3 transcatheter heart valve system. EuroIntervention 2019 Jan 20;14(13):1378-1385.

Martin MH, et al. Safety and feasibility of Melody transcatheter pulmonary valve replacement in the native right ventricular outflow tract: A multicenter pediatric heart network scholar study. JACC Cardiovasc Interv 2018 Aug 27;11(16):1642-1650.

*McElhinney DB, et al. Short- and medium-term outcomes after transcatheter pulmonary valve placement in the expanded multicenter US melody valve trial. Circulation 2010 Aug 3;122(5):507-16.

*McElhinney DB, et al. Infective endocarditis after transcatheter pulmonary valve replacement using the Melody valve: combined results of 3 prospective North American and European studies. Circ Cardiovasc Interv 2013 Jun;6(3):292-300.

McElhinney DB, et al. Multicenter study of endocarditis after transcatheter pulmonary valve replacement. J Am Coll Cardiol 2021;78:575-589.

Morgan GJ, et al. SAPIEN valve for percutaneous transcatheter pulmonary valve replacement without "pre-stenting": a multi-institutional experience. Catheter Cardiovasc Interv 2019;93(2):324-329.

National Institute for Health and Clinical Excellence (NICE). Interventional procedure guidance 436. Percutaneous pulmonary valve implantation for right ventricular outflow tract dysfunction. UK: NICE; 2013 January. [<http://www.nice.org.uk/guidance/ipg436>] Accessed 02/21/23.

Nordmeyer J, et al. Acute and midterm outcomes of the post-approval MELODY Registry: a multicentre registry of transcatheter pulmonary valve implantation. Eur Heart J 2019 Jul 14;40(27):2255-2264.

Plessis J, et al. Edwards SAPIEN transcatheter pulmonary valve implantation: results from a french registry. JACC Cardiovasc Interv 2018 Oct 8;11(19):1909-1916.

Ran L, et al. Percutaneous pulmonary valve implantation in patients with right ventricular outflow tract dysfunction: a systematic review and meta-analysis. Ther Adv Chronic Dis 2019 Jun 14;10:2040622319857635.

Ribeiro JM, et al. Transcatheter versus surgical pulmonary valve replacement: a systemic review and meta-analysis. Ann Thorac Surg 2020;110:1751-61.

Ruiz CE, et al. Are we ready for transcatheter pulmonary valve replacement in native right ventricular outflow tract? JACC Cardiovasc Interv 2018 Aug 27;11(16):1651-1653.

Sinha S, et al. Initial results from the off-label use of the SAPIEN S3 valve for percutaneous transcatheter pulmonary valve replacement: A multi-institutional experience. Catheter Cardiovasc Interv 2019 Feb 15;93(3):455-463.

*Stout KK, et al. 2018 AHA/ACC guideline for the management of adults with congenital heart disease: a report of the American College of Cardiology/American Heart Association Task Force on clinical practice guidelines. J Am Coll Cardiol 2019 Apr 2;73(12):e81-e192.

U.S. Food and Drug Administration Device Approvals and Clearances. Humanitarian Device Approval, Medtronic Melody® Transcatheter Pulmonary Valve - H080002.

[<https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfhde/hde.cfm?id=H080002>] Accessed 02/21/23.

Wadia SK, et al. Postoperative and short-term atrial tachyarrhythmia burdens after transcatheter vs surgical pulmonary valve replacement among congenital heartdisease patients. Congenit Heart Dis 2019;14(5):838-845.

*Warnes CA, et al. ACC/AHA 2008 Guidelines for the management of adults with congenital heart disease: Executive

Medical Policy: Transcatheter Heart Valve Procedures

Policy Number: 7.01.109

Page 16 of 17

Summary: a report of the American College of Cardiology/American Heart Association Task Force on practice guidelines. Circulation 2008 Dec 2;118(23):2395-451.

Tricuspid Valve:

Alperi AA, et al. Early and mid-term outcomes of transcatheter tricuspid valve repair: systematic review and meta-analysis of observational studies. Rev Esp Cardiol (Engl Ed) 2022 Jun 2:S1885-5857

Asmarats L, et al. Transcatheter tricuspid valve replacement for treating severe tricuspid regurgitation: initial experience with the NaviGate bioprosthesis. Can J Cardiol. 2018;34(10):1370.e1375-1370.e1377.

Curio J et al. Update on the current landscape of transcatheter options for tricuspid regurgitation treatment. Interv Cardiol 2019 May;14(2): 54–61.

Hahn RT, et al. Early single-site experience with transcatheter tricuspid valve replacement. JACC Cardiovasc Imaging 2019;Mar;12(3):416-429.

*Nickenig G, et al. 6-Month outcomes of tricuspid valve reconstruction for patients with severe tricuspid regurgitation. J Am Coll Cardiol 2019;73(15):1905.

*Nishimura RA, et al. 2017 AHA/ACC Focused update of the 2014 AHA/ACC guideline for the management of patients with valvular heart disease: a report of the American College of Cardiology/American Heart Association Task Force on clinical practice guidelines. Circulation 2017.

Ozam M et al. Transcatheter tricuspid valve replacement: Principles and Design. Front Cardiovasc Med 2018;5:129.

Ranocchi F1, et al. Minimally invasive implantation of a Sapien 3 Ultra Valve in a degenerated tricuspid bioprosthesis. J Card Surg 2020 Jun;35(6):1371-1374.

Ruf TF, et al. Short-term clinical outcomes of transcatheter tricuspid valve repair with the third generation MitraClip XTR System. J Am Coll Cardiol Intv Jun 2021;14(1):1231–40.

*Taramasso M, et al. Outcomes after current transcatheter tricuspid valve intervention: mid-term results from the International TriValve Registry. JACC Cardiovasc Interv 2019;Jan 28;12(2):155-165.

*Key Articles

KEY WORDS

Transcatheter aortic valve implantation (TAVI), Transcatheter aortic valve replacement (TAVR), Surgical aortic valve repair (SAVR), Transcatheter mitral valve repair (TMVR), MitraClip, Transcatheter pulmonic valve implantation (TPVI), Percutaneous pulmonary valve implantation (PPVI), Melody TPV System, Harmony TPV System, Transcatheter Tricuspid Valve Repair (TTVR), cerebral embolic protection device (CEPD)

CMS COVERAGE FOR MEDICARE PRODUCT MEMBERS

TAVR

Based upon our review, the Transcatheter Aortic Valve Replacement (TAVR) procedure is addressed in the following National Medicare coverage determination. [<https://www.cms.gov/medicare-coverage-database/details/ncd-details.aspx?NCDId=355>]

The Centers for Medicare & Medicaid Services (CMS) covers TAVR under Coverage with Evidence Development (CED) with certain conditions. Please refer to the following CED website for Medicare Members: [<https://www.cms.gov/Medicare/Coverage/Coverage-with-Evidence-Development/TAVR>]

There is currently a Decision Memo for Transcatheter Aortic Valve Replacement (TAVR)(CAG-00430R). Please refer to the following website: [<https://www.cms.gov/medicare-coverage-database/details/nca-decision-memo.aspx?NCAId=293&bc=ACAAAAAAQAAA&>]

TMVR

Medical Policy: Transcatheter Heart Valve Procedures

Policy Number: 7.01.109

Page 17 of 17

Based upon our review, the Transcatheter Mitral Valve Repair (TMVR) is addressed in the following National Medicare coverage determination. [<https://www.cms.gov/medicare-coverage-database/details/ncd-details.aspx?NCDId=363&ncdver=1&DocID=20.33&bc=gAAAAAgAAAA&>]

The Centers for Medicare & Medicaid Services (CMS) covers TMVR for MR under Coverage with Evidence Development (CED). Please refer to the following CED website for Medicare Members:

[<https://www.cms.gov/Medicare/Coverage/Coverage-with-Evidence-Development/TMVR>] _

There is currently a Decision Memo for Transcatheter Mitral Valve Repair (TMVR). Please refer to the following website: [<https://www.cms.gov/medicare-coverage-database/details/nca-decision-memo.aspx?NCAId=273>]

TPVI / TTVR

Based upon our review, Transcatheter Pulmonary Valve Implantation (TPVI) and Transcatheter Tricuspid Valve Repair or Replacement (TTVR) are not addressed in National or Regional Medicare coverage determinations or policies.