

MEDICAL POLICY



SUBJECT: DEEP BRAIN STIMULATION	EFFECTIVE DATE: 09/16/99 REVISED DATE: 07/19/01, 05/16/02, 03/20/03, 03/18/04, 03/17/05, 01/19/06, 01/18/07, 11/15/07, 11/20/08, 10/29/09, 10/28/10, 09/15/11, 08/16/12, 07/18/13, 06/19/14, 05/29/15, 06/16/16, 05/18/17, 04/19/18
POLICY NUMBER: 7.01.23 CATEGORY: Technology Assessment	PAGE: 1 OF: 16
<ul style="list-style-type: none">• <i>If a product excludes coverage for a service, it is not covered, and medical policy criteria do not apply.</i>• <i>If a commercial product, including an Essential Plan product, covers a specific service, medical policy criteria apply to the benefit.</i>• <i>If a Medicare product covers a specific service, and there is no national or local Medicare coverage decision for the service, medical policy criteria apply to the benefit.</i>	

POLICY STATEMENT:

- I. Based upon our criteria and assessment of the peer-reviewed literature, unilateral or bilateral deep brain stimulation of the ventral intermediate nucleus (VIM) thalamus has been medically proven to be effective and therefore **medically appropriate** for disabling, medically unresponsive essential tremor or tremor due to Parkinson’s disease (bilateral DBS would be utilized for bilateral tremor).

Disabling, medically unresponsive tremor is defined as both of the following:
 - A. tremor causes significant limitation in daily activities; and
 - B. inadequate control by maximal dosage of medication for at least 3 months before implant.
- II. Based upon our criteria and assessment of the peer-reviewed literature, unilateral or bilateral deep brain stimulation of the subthalamic nucleus (STN) or of the globus pallidus interna (GPi) has been medically proven to be effective and therefore **medically appropriate** for treatment of advanced Parkinson’s disease. All of the following criteria must be met:
 - A. the patient has a diagnosis of idiopathic (not secondary) Parkinson’s disease;
 - B. the patient’s Parkinson’s disease was previously responsive to levodopa therapy but is now medically intractable; and
 - C. the patient has severe levodopa-induced dyskinesia or disease characterized by severe bradykinesia, rigidity, tremor or dystonia or by marked “on-off” fluctuations.
- III. Based upon our criteria and assessment of peer-reviewed literature, bilateral deep brain stimulation of the subthalamic nucleus (STN) or of the globus pallidus interna (GPi) has been medically proven to be effective and therefore **medically appropriate** for treatment of patients who have had a Parkinson’s diagnosis for at least four years duration and who have recently developed motor complications that cause significant limitations in daily activities (patient need not be considered as having advanced Parkinson’s disease).
- IV. Based upon our criteria and assessment of peer-reviewed literature, unilateral or bilateral deep brain stimulation of the GPi or STN has been medically proven effective and therefore **medically appropriate** in patients 7 years of age or greater who experience chronic, intractable, primary dystonia, including generalized and focal dystonia.
- V. Based upon our criteria and assessment of the peer-reviewed literature, there is insufficient clinical evidence to support the safety and efficacy of deep brain stimulation, and is therefore considered **investigational** for the following conditions, including but not limited to:
 - A. Multiple Sclerosis,
 - B. post-traumatic dyskinesia;
 - C. all other movement disorders;
 - D. chronic pain syndromes, including cluster headache;
 - E. tardive dyskinesia;
 - F. epilepsy;
 - G. Tourette syndrome;
 - H. Dementias, including Alzheimer’s disease;
 - I. Eating disorders, including Anorexia nervosa;

<p>SUBJECT: DEEP BRAIN STIMULATION</p> <p>POLICY NUMBER: 7.01.23</p> <p>CATEGORY: Technology Assessment</p>	<p>EFFECTIVE DATE: 09/16/99</p> <p>REVISED DATE: 07/19/01, 05/16/02, 03/20/03, 03/18/04, 03/17/05, 01/19/06, 01/18/07, 11/15/07, 11/20/08, 10/29/09, 10/28/10, 09/15/11, 08/16/12, 07/18/13, 06/19/14, 05/29/15, 06/16/16, 05/18/17, 04/19/18</p> <p>PAGE: 2 OF: 16</p>
--	--

- J. Alcohol addiction;
- K. treatment-resistant depression; or
- L. treatment-resistant obsessive compulsive disorder.

VI. Contraindications to deep brain stimulation include:

- A. patients who are not good surgical risks because of unstable medical problems;
- B. patients who have a cardiac pacemaker;
- C. patients who have medical conditions that require repeated MRI; or
- D. patients who have neuropsychiatric disease that may interfere with their ability to benefit from deep brain stimulation.

This medical policy does not address occipital nerve stimulation for chronic migraines or occipital neuralgia. In occipital nerve stimulation the neurostimulator delivers electrical impulses via insulated lead wires tunneled under the skin near the occipital nerves at the base of the head.

This medical policy does not address stimulation of the motor cortex, which has been investigated as a treatment for patients with chronic, refractory neuropathic pain and extremity weakness due to stroke. In motor cortex stimulation, electrodes are implanted subdurally over the sensorimotor cortex.

POLICY GUIDELINES:

- I. Bilateral stimulators may be implanted simultaneously or in staged procedures.
- II. The Federal Employee Health Benefit Program (FEHBP/FEP) requires that procedures, devices or laboratory tests approved by the U.S. Food and Drug Administration (FDA) may not be considered investigational and thus these procedures, devices or laboratory tests may be assessed only on the basis of their medical necessity.

DESCRIPTION:

Deep brain stimulation (DBS) has been investigated as an alternative to permanent neuroablative procedures such as thalamotomy and pallidotomy. The procedure involves the stereotactic placement of an electrode into a targeted region of the brain. The electrode is then attached, via a cable/wire, to a programmable stimulator implanted subcutaneously. Deep brain stimulation is designed to turn off overactive brain regions without destroying them. The immediate advantage of DBS over conventional destructive surgery is that the lesions are titratable and hence reversible. After implantation, noninvasive programming of the neurostimulator can be adjusted to the patient's symptoms.

The effect of DBS depends on where the electrodes are placed. The 3 common target sites are the VIM thalamus, globulus pallidus interna and subthalamic nucleus. Whereas unilateral/bilateral DBS of the thalamus is utilized to treat essential tremor or tremors of advanced Parkinson's disease, bilateral deep brain stimulation of the subthalamic nucleus (STN) or of the globus pallidus interna (GPi) is used for treatment of the entire constellation of Parkinsonian symptoms (e.g., tremor, rigidity, and bradykinesia). Deep brain stimulation is performed at specialty centers.

DBS has also been investigated for the treatment of primary dystonia, defined as a neurological movement disorder characterized by involuntary and painful muscle contractions and contortions. Dystonia can be classified according to cause and the bodily distribution of symptoms. Primary or idiopathic dystonia is not associated with any other pathology whereas, secondary dystonia is caused by a known insult (e.g., trauma, infarct, stroke) to the basal ganglia. Generalized dystonia affects a wide range of body areas and focal dystonia affects specific body parts (e.g., spasmodic torticollis/cervical dystonia, blepharospasm). Dystonia is the third most common movement disorder, behind Parkinson's disease and essential tremor. Unless contraindicated, DBS of either the GPi or STN requires a bilateral procedure.

In addition to essential tremors, Parkinson's disease, and dystonia, deep brain stimulation is also being investigated for disorders such as major depression, cluster headaches, chronic pain syndromes, Tourette syndrome, epilepsy and obsessive-compulsive disorder.

<p>SUBJECT: DEEP BRAIN STIMULATION</p> <p>POLICY NUMBER: 7.01.23</p> <p>CATEGORY: Technology Assessment</p>	<p>EFFECTIVE DATE: 09/16/99</p> <p>REVISED DATE: 07/19/01, 05/16/02, 03/20/03, 03/18/04, 03/17/05, 01/19/06, 01/18/07, 11/15/07, 11/20/08, 10/29/09, 10/28/10, 09/15/11, 08/16/12, 07/18/13, 06/19/14, 05/29/15, 06/16/16, 05/18/17, 04/19/18</p> <p>PAGE: 3 OF: 16</p>
--	--

Directional DBS

Conventional DBS systems use ring-shaped electrodes, which generate an approximately spherical electrical field. In these systems, programming of polarity and stimulation pulse parameters allows only limited control of the shape of the volume of tissue activated. While physicians try to target a very specific area of the brain, with conventional DBS there is a risk of stimulating neighboring regions as they can't steer the stimulation precisely. Directional DBS systems use novel lead designs with segmented, multi-contact electrodes that allow for the activation of individual electrode contacts and also allows the physician to specify the exact amount of current needed for every contact of the electrode. Through activating specific electrode contacts and defining the amount of stimulation for each contact, stimulation precision is significantly increased. More precise stimulation is thought to reduce side effects of DBS such as muscle contractions, dysarthria and cognitive or behavioral disturbances sometimes seen in conventional DBS.

RATIONALE:

The FDA approved the Activa® Tremor Control System (Medtronic Inc.) for DBS. While the original 1997 FDA-labeled indications were limited to unilateral implantation of the device for the treatment of tremor, in January 2002, the FDA-labeled indications were expanded to include bilateral implantation as a treatment to decrease the symptoms of advanced PD that are not controlled by medication. In February 2016, the FDA expanded the approval for Medtronic's DBS for Parkinson's disease. The expanded approval covers patients who have had a Parkinson's diagnosis for four years and who have recently developed motor complications, or have long standing motor complications that cannot be controlled with drugs. The expanded approval is based on data from the EARLYSTIM clinical study (Schuepbach WM, et al. 2013) which found that patients treated with Medtronic DBS Therapy and best medical therapy (BMT) reported a mean improvement of 26 percent in their disease-related quality of life at two years, compared to a one percent decline in patients treated with BMT alone. In a study of patients with longer-standing motor complications, DBS patients' quality of life improved 20 percent from baseline to six months compared to no improvement in the patients treated with BMT alone.

In April of 2003, the FDA gave Humanitarian Device Exemption (HDE) approval to the Activa Therapy system for the unilateral or bilateral stimulation of the internal GPi or STN to aid in the management of chronic, intractable (drug resistant) primary dystonia, including generalized and/or segmental dystonia, hemidystonia and cervical dystonia in patients seven years of age or greater.

The Brio Neuromodulation System (St. Jude Medical) received FDA approval in June of 2015. The device is indicated for the following conditions: 1) bilateral stimulation of the subthalamic nucleus (STN) as an adjunctive therapy to reduce some of the symptoms of advanced levodopa-responsive Parkinson's disease that are not adequately controlled by medications; and 2) unilateral or bilateral stimulation of the ventral intermediate nucleus (VIM) of the thalamus for the suppression of disabling upper extremity tremor in adult essential tremor patients whose tremor is not adequately controlled by medications and where the tremor constitutes a significant functional disability. The Brio device differs from the Activa system in how it provides stimulation to the brain- using a constant current of electricity to the brain versus constant voltage. Per the FDA Summary of Safety and Effectiveness Data: Data supporting its use come from two clinical trials of the device, one in 136 PD patients and the other in 127 patients with ET. In both studies, symptoms were not adequately controlled with medication. The system was used as an adjunct to medication for the patients with Parkinson's, while "the majority of patients with essential tremor who used the device were able to control their symptoms without the need for medications," the FDA said. All patients in the studies were implanted with the system; PD patients were evaluated at 3 months, and the ET patients after 6 months of therapy. "Both groups showed statistically significant improvement on their primary effectiveness endpoint when the device was turned on compared to when it was turned off," the statement notes.

Published clinical trials have provided evidence to support the efficacy and safety of unilateral deep brain stimulation of the VIM thalamus for essential tremor and for tremor of Parkinson's disease and of bilateral deep brain stimulation of the STN or GPi for advanced Parkinson's disease. In studies of unilateral thalamic DBS, tremor suppression was either total or clinically significant in 82-91% of patients who underwent implantation. Results were durable and side effects were minimal. An additional benefit of DBS is that recurrence of tremor may be managed by changes in stimulation parameters.

<p>SUBJECT: DEEP BRAIN STIMULATION</p> <p>POLICY NUMBER: 7.01.23</p> <p>CATEGORY: Technology Assessment</p>	<p>EFFECTIVE DATE: 09/16/99</p> <p>REVISED DATE: 07/19/01, 05/16/02, 03/20/03, 03/18/04, 03/17/05, 01/19/06, 01/18/07, 11/15/07, 11/20/08, 10/29/09, 10/28/10, 09/15/11, 08/16/12, 07/18/13, 06/19/14, 05/29/15, 06/16/16, 05/18/17, 04/19/18</p> <p>PAGE: 4 OF: 16</p>
--	--

Although long-term data are minimal, studies have demonstrated that bilateral stimulation of the GPi or STN results in improvements of neurologic function. Case series investigating the use of DBS for the treatment of dystonia found that patients with primary dystonia experienced significant improvement in movement and in ADL's, but those patients with secondary dystonia experienced little improvement.

Directional DBS

The St. Jude Medical Infinity™ DBS system is the first FDA approved system to feature a directional lead, designed to deliver electrical current to a specific target in the brain and therefore, proposed to minimize unwanted side effects from brain stimulation to non-targeted areas. On September 19, 2016, the St. Jude Medical Infinity™ DBS system was FDA approved as a supplement to an earlier Premarket Approval (PMA) for the St. Jude Medical Brio Neurostimulation system. This approval was for a change in design, components, specifications, and material. The most recent supplemental approval pertaining to the Infinity neurostimulation system was issued on May 1, 2017, and was for a firmware update. According to the manufacturer, the Infinity DBS System is indicated for “bilateral stimulation of the subthalamic nucleus (STN) as an adjunctive therapy to reduce some of the symptoms of advanced levodopa-responsive Parkinson’s disease that are not adequately controlled by medications, and unilateral or bilateral stimulation of the ventral intermediate nucleus (VIM) of the thalamus for the suppression of disabling upper extremity tremor in adult essential tremor patients whose tremor is not adequately controlled by medications and where the tremor constitutes a significant functional disability”.

In December 2017, Boston Scientific Inc. received FDA approval (PMA) for its Vercise Deep Brain Stimulation System that includes directional lead technology. The Vercise Deep Brain Stimulation (DBS) System is indicated for use in bilateral stimulation of the subthalamic nucleus (STN) as an adjunctive therapy in reducing some of the symptoms of moderate to advanced levodopa-responsive Parkinson’s disease (PD) that are not adequately controlled with medication. The Vercise DBS System utilizes current steering across eight contacts per DBS Lead, which is intended to provide precise positioning of stimulation.

The FDA approved Medtronic’s *ReClaim* Deep Brain Stimulator device as the first implant to treat severe obsessive-compulsive disorder under a HDE approval in February 2009. The device is indicated for bilateral stimulation of the anterior limb of the internal capsule (AIC) as an adjunct to medications and as an alternative to anterior capsulotomy for the treatment of chronic, severe, treatment-resistant obsessive compulsive disorder in adult patients who have failed at least three selective serotonin reuptake inhibitors (SSRIs). The approval of the human device exemption was based on a review of data from 26 patients with severe treatment resistant OCD who were treated with the device at four sites. On average, patients had a 40 percent reduction in their symptoms after 12 months of therapy. One of the major limitations of this study was the fact that many of the study population were aware of when the device was turned on and off, so investigators were unable to rule out that some of the improvements were due to a placebo effect. While there is limited evidence to suggest that DBS may be an option for patients with severe, disabling OCD, well designed studies are necessary to demonstrate its long term safety and efficacy.

Published clinical trials have not provided evidence to support the efficacy and safety of deep brain stimulation for conditions including, but not limited to, Multiple Sclerosis, post-traumatic dyskinesia, treatment-resistant depression, Alzheimer’s disease, Tourette syndrome, or for bilateral deep brain stimulation of the VIM thalamus. Studies of deep brain stimulation for the treatment of chronic pain have not provided evidence that this is an effective treatment method over already established treatment methods.

Results of Medtronic’s SANTE trial (Fisher, et al. 2010) show promising outcomes on the adjunct use of deep brain stimulation of the anterior nucleus of the thalamus over placebo stimulation for patients suffering from severe, refractory, partial-onset seizures. Two years after implantation of the device, seizures were reduced by a median 56% compared with baseline and 14 patients (12.7%) became seizure-free for at least 6 months. The FDA has not yet granted FDA approval for the use of DBS in epilepsy and longer-term studies are needed to better define its safety, efficacy and the subset of patients who would benefit most from this treatment.

SUBJECT: DEEP BRAIN STIMULATION POLICY NUMBER: 7.01.23 CATEGORY: Technology Assessment	EFFECTIVE DATE: 09/16/99 REVISED DATE: 07/19/01, 05/16/02, 03/20/03, 03/18/04, 03/17/05, 01/19/06, 01/18/07, 11/15/07, 11/20/08, 10/29/09, 10/28/10, 09/15/11, 08/16/12, 07/18/13, 06/19/14, 05/29/15, 06/16/16, 05/18/17, 04/19/18 PAGE: 5 OF: 16
---	---

CODES: Number Description

Eligibility for reimbursement is based upon the benefits set forth in the member's subscriber contract.

CODES MAY NOT BE COVERED UNDER ALL CIRCUMSTANCES. PLEASE READ THE POLICY AND GUIDELINES STATEMENTS CAREFULLY.

Codes may not be all inclusive as the AMA and CMS code updates may occur more frequently than policy updates.

CPT:	61850	Twist drill or burr hole(s) for implantation of neurostimulator electrodes, cortical
	61863	Twist drill, burr hole, craniotomy or craniectomy with stereotactic implantation of neurostimulator electrode array in subcortical site (eg, thalamus, globus pallidus, subthalamic nucleus, periventricular, periaqueductal gray), without use of intraoperative microelectrode recording; first array
	61864	each additional array
	61867	Twist drill, burr hole, craniotomy or craniectomy with stereotactic implantation of neurostimulator electrode array in subcortical site (eg, thalamus, globus pallidus, subthalamic nucleus, periventricular, periaqueductal gray), with use of intraoperative microelectrode recording; first array
	61868	each additional array
	61880	Revision or removal of intracranial neurostimulator electrodes
	61885	Insertion or replacement of cranial neurostimulator pulse generator or receiver, direct or inductive coupling; with connection to a single electrode array
	61886	with connection to two or more electrode arrays
	61888	revision or removal of cranial neurostimulator pulse generator or receiver
	95970	electronic analysis of implanted neurostimulator pulse generator system; simple or complex brain, spinal cord, or peripheral neurostimulator pulse generator/transmitter, without programming
	95978	Electronic analysis of implanted neurostimulator pulse generator system (eg, rate, pulse amplitude and duration, battery status, electrode selectability and polarity, impedance and patient compliance measurements) complex deep brain stimulator pulse generator/transmitter, with initial or subsequent programming, first hour
	95979	each additional 30 minutes after first hour

Copyright © 2018 American Medical Association, Chicago, IL

HCPCS:	C1767	Generator, neurostimulator (implantable), nonrechargeable
	C1820	Generator, neurostimulator (implantable), non high frequency with rechargeable battery and charging system
	C1822	Generator, neurostimulator (implantable), high frequency with rechargeable battery and charging system
	C1787	Patient programmer; neurostimulator
	L8679	Implantable neurostimulator pulse generator, any type
	L8680	Implantable neurostimulator electrode, each

<p>SUBJECT: DEEP BRAIN STIMULATION</p> <p>POLICY NUMBER: 7.01.23</p> <p>CATEGORY: Technology Assessment</p>	<p>EFFECTIVE DATE: 09/16/99</p> <p>REVISED DATE: 07/19/01, 05/16/02, 03/20/03, 03/18/04, 03/17/05, 01/19/06, 01/18/07, 11/15/07, 11/20/08, 10/29/09, 10/28/10, 09/15/11, 08/16/12, 07/18/13, 06/19/14, 05/29/15, 06/16/16, 05/18/17, 04/19/18</p> <p>PAGE: 6 OF: 16</p>
--	--

- L8681 Patient programmer (external) for use with implantable programmable neurostimulator pulse generator
- L8682 Implantable neurostimulator radiofrequency receiver
- L8683 Radiofrequency transmitter (external) for use with implantable neurostimulator radiofrequency receiver
- L8685 Implantable neurostimulator pulse generator, single array, rechargeable, includes extension
- L8686 Implantable neurostimulator pulse generator, single array, non-rechargeable, includes extension
- L8687 Implantable neurostimulator pulse generator, dual array, rechargeable, includes extension
- L8688 Implantable neurostimulator pulse generator, dual array, non-rechargeable, includes extension
- L8689 External recharging system for battery (internal) for use with implantable neurostimulator

ICD 10:

- G20 Parkinson's disease
- G21.11-G21.9 Parkinsonism (code range)
- G24.02-G24.3 Dystonia (code range)
- G24.5 Blepharospasm
- G24.8 Other dystonia
- G25.0 Essential tremor
- G25.1 Drug-induced tremor
- G25.2 Other specified forms of tremor
- G80.3 Athetoid cerebral palsy

REFERENCES:

*Abelson JL, et al. Deep brain stimulation for refractory obsessive-compulsive disorder. Biol Psych 2005 Mar 1;57(5):510-6.

*Ackermans L, et al. Double-blind clinical trial of thalamic stimulation in patients with Tourette syndrome. Brain 2011 March;134(Pt 3):832-44.

*Air EL, et al. Deep brain stimulation in children: experience and technical pearls. J Neurosurg Pediatr 2011 Dec;8(6):566-74.

Akram H, et al. Optimal deep brain stimulation site and target connectivity for chronic cluster headache. Neurology 2017 Nov 14;89(20):2083-2091.

Alonso P, et al. Deep brain stimulation for obsessive-compulsive disorder: a meta-analysis of treatment options and predictors of response. PLoS One 2015 Jul 24;10(7):e0133591.

Altinay M, et al. A comprehensive review of the use of deep brain stimulation (DBS) in treatment of psychiatric and headache disorders. Headache 2015 Feb;55(2):345-50.

<p>SUBJECT: DEEP BRAIN STIMULATION</p> <p>POLICY NUMBER: 7.01.23</p> <p>CATEGORY: Technology Assessment</p>	<p>EFFECTIVE DATE: 09/16/99</p> <p>REVISED DATE: 07/19/01, 05/16/02, 03/20/03, 03/18/04, 03/17/05, 01/19/06, 01/18/07, 11/15/07, 11/20/08, 10/29/09, 10/28/10, 09/15/11, 08/16/12, 07/18/13, 06/19/14, 05/29/15, 06/16/16, 05/18/17, 04/19/18</p> <p>PAGE: 7 OF: 16</p>
--	--

American Society of Stereotactic and Functional Neurosurgery. Deep brain stimulation: indications, techniques and practice parameters. 2010 [http://www.assfn.org/practice/dbs_statement.pdf] accessed 3/13/18.

*Anderson VC, et al. Pallidal vs subthalamic nucleus deep brain stimulation in Parkinson's disease. Arch Neurol 2005 Apr;62(4):554-60.

*Andrade P, et al. A systematic review of the efficacy of globus pallidus stimulation in the treatment of Parkinson's disease. J Clin Neurosci 2009 Jul;16(7):877-81.

*Andrade P, et al. Neurostimulatory and ablative treatment options in major depressive disorder: a systematic review. Acta Neurochir 2010 Apr;152(4):566-77.

*Antonini A, et al. A 5-year prospective assessment of advanced Parkinson disease patients treated with subcutaneous apomorphine infusion or deep brain stimulation. J Neurol 2011 Apr;258(4):579-85.

*Appleby BS, et al. Psychiatric and neuropsychiatric adverse events associated with deep brain stimulation: A meta-analysis of ten years' experience. Mov Disord 2007 Sep 15;22(12):1722-8.

Aviles-Olmos I, et al. Long-term outcome of subthalamic nucleus deep brain stimulation for Parkinson's disease using an MRI-guided and MRI-verified approach. J Neurol Neurosurg Psychiatry 2014 Dec;85(12):1419-25.

*Baumer T, et al. Effects of DBS, premotor rTMS, and levodopa on motor function and silent period in advance Parkinson's disease. Mov Disord 2009 Apr 15;24(5):672-6.

Bergfeld IO, et al. Deep brain stimulation of the ventral anterior limb of the internal capsule for treatment-resistant depression: a randomized clinical trial. JAMA Psychiatry May 1 2016;73(5):456-464.

Berlim MT, et al. Effectiveness and acceptability of deep brain stimulation (DBS) of the subgenual cingulate cortex for treatment-resistant depression: a systematic review and exploratory meta-analysis. J Affective Disord 2014 Apr;159:31-8.

*Bewernick BH, et al. Nucleus accumbens deep brain stimulation decreases ratings of depression and anxiety in treatment-resistant depression. Biol Psychiatry 2010 Jan 15;67(2):110-6.

Bhidayasiri R, et al. Evidence-based guideline: treatment of tardive syndromes: report of the Guideline Development Subcommittee of the American Academy of Neurology. Neurology 2013 Jul 30;81(5):463-9.

*Blomstedt P, et al. Deep brain stimulation in the treatment of depression. Acta Psychiatr Scand 2011 Jan;123(1):4-11.

Blomstedt P, et al. Deep brain stimulation in the treatment of obsessive-compulsive disorder. World Neurosurg 2013 Dec;80(6):e245-53.

*BlueCross BlueShield Association Technology Evaluation Center (TEC). Bilateral DBS of the subthalamic nucleus or the globus pallidus interna for treatment of advanced Parkinson's disease. 2002 Feb.

BlueCross BlueShield Association. Deep brain stimulation. Medical Policy Reference Manual Policy #7.01.63. 2017 May 8.

Boccard SG, et al. Long-term outcomes of deep brain stimulation for neuropathic pain. Neurosurgery 2013 Feb;72(2):221-30.

Boccard SGJ, et al. Long-term results of deep brain stimulation of the anterior cingulate cortex for neuropathic pain. World Neurosurg 2017 Oct;106:625-637.

*Bronstein JM, et al. Deep brain stimulation for Parkinson disease: an expert consensus and review of key issues. Arch Neurol 2011 Feb;68(2):165-71.

Bour LJ, et al. Directional recording of subthalamic spectral power densities in Parkinson's disease and the effect of steering deep brain stimulation. Brain Stimul 2015 July- Aug;8(4):730-741.

SUBJECT: DEEP BRAIN STIMULATION POLICY NUMBER: 7.01.23 CATEGORY: Technology Assessment	EFFECTIVE DATE: 09/16/99 REVISED DATE: 07/19/01, 05/16/02, 03/20/03, 03/18/04, 03/17/05, 01/19/06, 01/18/07, 11/15/07, 11/20/08, 10/29/09, 10/28/10, 09/15/11, 08/16/12, 07/18/13, 06/19/14, 05/29/15, 06/16/16, 05/18/17, 04/19/18 PAGE: 8 OF: 16
---	---

- *Cannon E, et al. Deep brain stimulation of anteromedial globus pallidus interna for severe Tourette's syndrome. Am J Psychiatry 2012 Aug 1;169(8):860-6.
- *Capecci M, et al. Functional improvement after subthalamic stimulation in Parkinson's disease: a non-equivalent controlled study with 12-24 month follow up. J Neurol Neurosurg Psych 2005 Jun;76(6):769-74.
- *Castrioto A, et al. Ten-year outcome of subthalamic stimulation in Parkinson disease: a blinded evaluation. Arch Neurol 2011 Dec;68(12):1550-6.
- *Charles PD, et al. Is deep brain stimulation neuroprotective if applied early in the course of PD? Nat Clin Pract Neurol 2008 Aug;4(8):424-6.
- Chiou SM, et al. One-year outcome of bilateral subthalamic stimulation in Parkinson disease: an Eastern experience. World Neurosurg 2015 Nov;84(5):1294-8.
- Chiou SM. Benefits of subthalamic stimulation for elderly parkinsonian patients aged 70 years or older. Clin Neurol Neurosurg 2016 Oct;149:81-86.
- *Clarke CE, et al. Systematic review of apomorphine, levodopa infusion and deep brain stimulation in advanced Parkinson's disease. Parkinsonism Relat Disord 2009 Dec;15(10):728-41.
- Combs HL, et al. Cognition and depression following deep brain stimulation of the subthalamic nucleus and globus pallidus par internus in Parkinson's disease: A meta-analysis. Neuropsychol Rev 2015 Dec;25(4):439-54.
- Contarino MF, et al. Directional steering: a novel approach to deep brain stimulation. Neurology 2014 Sept 23;83(13):1163-1169.
- *Coubes, P, et al. Treatment of DYT11-generalized dystonia by stimulation of the internal globus pallidus. Lancet 2000;355:2220-1.
- *Coubes P, et al. Electrical stimulation of the globus pallidus internus in patients with primary generalized dystonia: long-term results. J Neurosurg 2004 Aug;101(2):189-94.
- Cury RG, et al. Effects of deep brain stimulation on pain and other nonmotor symptoms in Parkinson disease. Neurology 2014 Oct 14;83(16):1403-9.
- *Damier P, et al. Bilateral deep brain stimulation of the globus pallidus to treat tardive dyskinesia. Arch Gen Psych 2007 Feb;64(2):170-6.
- Dembek TA, et al. Directional DBS increases side-effect thresholds- a prospective, double-blind trial. Mov Disord 2017 Oct;32(10):1380-1388.
- *Denys D, et al. Deep brain stimulation of the nucleus accumbens for treatment-refractory obsessive-compulsive disorder. Arch Gen Psychiatry 2010 Oct;67(10):1061-8.
- deSouza RM, et al. Timing of deep brain stimulation in Parkinson disease: a need for reappraisal? Ann Neurol 2013 May;73(5):565-75.
- *Deuschl G, et al. A randomized trial of deep-brain stimulation for Parkinson's disease. NEJM 2006 Aug 31;355(9):896-908.
- *Diamond A, et al. Globus pallidus deep brain stimulation in dystonia. Mov Disord 2006 May;21(5):692-5.
- Dougherty DD, et al. A randomized sham-controlled trial of deep brain stimulation of the ventral capsule/ventral striatum for chronic treatment-resistant depression. Biol Psychiatry 2015 Aug 15;78(4):240-8.
- *Egidi M, et al. A survey of Italian cases of dystonia treated by deep brain stimulation. J Neurosurg Sci 2007 Dec;51(4):153-8.

<p>SUBJECT: DEEP BRAIN STIMULATION</p> <p>POLICY NUMBER: 7.01.23</p> <p>CATEGORY: Technology Assessment</p>	<p>EFFECTIVE DATE: 09/16/99</p> <p>REVISED DATE: 07/19/01, 05/16/02, 03/20/03, 03/18/04, 03/17/05, 01/19/06, 01/18/07, 11/15/07, 11/20/08, 10/29/09, 10/28/10, 09/15/11, 08/16/12, 07/18/13, 06/19/14, 05/29/15, 06/16/16, 05/18/17, 04/19/18</p> <p>PAGE: 9 OF: 16</p>
--	--

Elias GJB, et al. Deep brain stimulation for stroke: current uses and future directions. Brain Stimul 2018 Jan-Feb;11(1):3-28.

*Eltahawy HA, et al. Primary dystonia is more responsive than secondary dystonia to pallidal interventions: outcomes after pallidotomy or pallidal deep brain stimulation. Neurosurg 2004 Mar;54(3):613-21.

*Esselink RA, et al. Unilateral pallidotomy versus bilateral subthalamic nucleus stimulation in PD: a randomized trial. Neurol 2004 Jan 27;62(2):201-7.

*Esselink RA, et al. Long-term superiority of subthalamic nucleus stimulation over pallidotomy in Parkinson disease. Neurology 2009 Jul 14;73(2):151-3.

*Figueiras-Mendez R, et al. Further supporting evidence of beneficial subthalamic stimulation in Parkinson's patients. Neurol 2002 Feb 12;58(3):469-70.

*Fisher R, et al. Electrical stimulation of the anterior nucleus of the thalamus for treatment of refractory epilepsy. Epilepsia 2010 May 5;51(5):899-908.

FitzGerald JJ, et al. Long-term outcome of deep brain stimulation in generalized dystonia: a series of 60 cases. J Neurol Neurosurg Psychiatry 2014 Dec;85(12):1371-6.

Floden D, et al. Predicting quality of life outcomes after subthalamic nucleus deep brain stimulation. Neurology 2014 Oct 28;83(18):1627-33.

*Fontaine D, et al. Anatomical location of effective deep brain stimulation electrodes in chronic cluster headache. Brain 2010 Apr;133(pt 4):1214-33.

*Ford B, et al. Subthalamic nucleus stimulation in advanced Parkinson's disease: blinded assessments at one year follow up. J Neurol Neurosurg Psych 2004 Sep;75(9):1255-9.

Fox MD, et al. Brain stimulation for torsion dystonia. JAMA Neurol 2015 Jun;72(6):713-9.

Fraint A, et al. Deep brain Stimulation in Tourette's syndrome. Front Neurol 2015 Aug 4;6:170.

*Fraix V, et al. Clinical and economic results of bilateral subthalamic nucleus stimulation in Parkinson's disease. J Neurol Neurosurg Psych 2006 Apr;77(4):443-9.

*Franzini A, et al. Stimulation of the posterior hypothalamus for treatment of chronic intractable cluster headaches: first reported series. Neurosurg 2003 May;52(5):1095-9.

Franzini A, et al. Deep-brain stimulation for aggressive and disruptive behavior. World Neurosurg 2013 Sep-Oct;80(3-4):S29.e11-14.

Fukaya C, et al. Predictive factors for long-term outcome of subthalamic nucleus deep brain stimulation for Parkinson's disease. Neurol Med Chir 2017 April 15;57(4):166-177.

*Germano IM, et al. Unilateral stimulation of the subthalamic nucleus in Parkinson disease: a double-blind 12-month evaluation study. J Neurosurg 2004 Jul;101(1):36-42.

*Gervais-Bernard H, et al. Bilateral subthalamic nucleus stimulation in advanced Parkinson's disease: five year follow-up. J Neurol 2009 Feb;256(2):225-33.

Gimeno H, et al. Improvement in upper limb function in children with dystonia following deep brain stimulation. Eur J Paediatr Neurol 2013 Jul;17(4):353-60.

Golestanirad L, et al. Efficacy and safety of pedunculopontine nuclei (PPN) deep brain stimulation in the treatment of gait disorders: a meta-analysis of clinical studies. Can J Neurol Sci 2016 Jan;43(1):120-126.

<p>SUBJECT: DEEP BRAIN STIMULATION</p> <p>POLICY NUMBER: 7.01.23</p> <p>CATEGORY: Technology Assessment</p>	<p>EFFECTIVE DATE: 09/16/99</p> <p>REVISED DATE: 07/19/01, 05/16/02, 03/20/03, 03/18/04, 03/17/05, 01/19/06, 01/18/07, 11/15/07, 11/20/08, 10/29/09, 10/28/10, 09/15/11, 08/16/12, 07/18/13, 06/19/14, 05/29/15, 06/16/16, 05/18/17, 04/19/18</p> <p>PAGE: 10 OF: 16</p>
--	---

- *Goodman WK, et al. Deep brain stimulation for intractable obsessive compulsive disorder: pilot study using a blinded, staggered-onset design. Biol Psychiatry 2010 Mar 15;67(6):535-42.
- *Greenberg BD, et al. Three-year outcomes in deep brain stimulation for highly resistant obsessive-compulsive disorder. Neuropsychopharmacol 2006 Nov;31(11):2384-93.
- *Greenberg BD, et al. Deep brain stimulation of the ventral internal capsule/ventral striatum for obsessive-compulsive disorder: worldwide experience. Mol Psychiatry 2008 May 20 [Epub ahead of print].
- *Grover PJ, et al. Deep brain stimulation for cluster headache. J Clin Neurosci 2009 Jul;16(7):861-6.
- *Gruber D, et al. Long-term effects of pallidal deep brain stimulation in tardive dystonia. Neurol 2009 Jul 7;73(1):53-8.
- *Halbig TD, et al. Pallidal stimulation in dystonia: effects on cognition, mood, and quality of life. J Neurol Neurosurg Psych 2005 Dec;76(12):1713-6.
- Hamani C, et al. Deep brain stimulation for obsessive-compulsive disorder: systematic review and evidence-based guideline sponsored by the American Society for Stereotactic and Functional Neurosurgery and the Congress of Neurological Surgeons (CNS) and endorsed by the CNS and American Association of Neurological Surgeons. Neurosurgery 2014 Oct;75(4):327-33.
- *Harries AM, et al. Deep brain stimulation of the subthalamic nucleus for advanced Parkinson disease using general anesthesia: long-term results. J Neurosurg 2012 Jan;116(1):107-13.
- *Hauptman JS, et al. Potential surgical targets for deep brain stimulation in treatment-resistant depression. Neurosurg Focus 2008;25(1):E3.
- *Haynes WI, et al. High-frequency stimulation of deep brain structures in obsessive-compulsive disorder: the search for a valid circuit. Eur J Neurosci 2010 Oct;32(7):1118-27.
- Hescham S, et al. Deep brain stimulation in dementia-related disorders. Neurosci Biobehav Rev 2013 Dec;37(10 Pt 2):2666-75.
- Holtzheimer PE, et al. Subcallosal cingulate deep brain stimulation for treatment-resistant unipolar and bipolar depression. Arch Gen Psychiatry 2012 Feb;69(2):150-8.
- *Hung SW, et al. Long-term outcome of bilateral pallidal deep brain stimulation for primary cervical dystonia. Neurol 2007 Feb 6;68(6):457-9.
- *Institute for Clinical Systems Improvement (ICSI). Technology Assessment Report. Deep brain stimulation for essential tremor and Parkinson's disease. Updated Apr 2005 [<http://icis.org/knowledge/detail.asp?catID=306&itemID=276>] accessed 3/13/18.
- *Kahn E, et al. Deep brain stimulation in early stage Parkinson's disease: operative experience from a prospective randomized clinical trial. J Neurol Neurosurg Psychiatry 2012 Feb;83(2):164-70.
- Kaminsska M, et al. Complications of deep brain stimulation (DBS) for dystonia in children- the challenges and 10-year experience in a large paediatric cohort. Eur J Paediatr Neurol 2017 Jan;21(1):168-175.
- Kefalopoulou Z, et al. Bilateral globus pallidus stimulation for severe Tourette's syndrome: a double-blind, randomized crossover trial. Lancet Neurol 2015 Jun;14(6):595-605.
- *Kennedy SH, et al. Canadian Network for Mood and Anxiety Treatments (CANMAT) Clinical guidelines for the management of major depressive disorder in adults. IV. Neurostimulation therapies. J Affect Disord 2009 Oct;117(Suppl 1):S44-53.
- *Kennedy SH, et al. Deep brain stimulation for treatment-resistant depression: follow-up after 3 to 6 years. Am J Psychiatry 2011 May;168(5):502-10.

SUBJECT: DEEP BRAIN STIMULATION POLICY NUMBER: 7.01.23 CATEGORY: Technology Assessment	EFFECTIVE DATE: 09/16/99 REVISED DATE: 07/19/01, 05/16/02, 03/20/03, 03/18/04, 03/17/05, 01/19/06, 01/18/07, 11/15/07, 11/20/08, 10/29/09, 10/28/10, 09/15/11, 08/16/12, 07/18/13, 06/19/14, 05/29/15, 06/16/16, 05/18/17, 04/19/18 PAGE: 11 OF: 16
---	--

Kim JH, et al. Effect of subthalamic deep brain stimulation on levodopa-induced dyskinesia in Parkinson's disease. Yonsei Med J 2015 Sep;56(5):1316-21.

Kisley S, et al. Deep brain stimulation for obsessive-compulsive disorder: a systematic review and meta-analysis. Psychol Med 2014 Dec;44(16):3533-42.

*Kiss ZH, et al. The Canadian multicentre study of deep brain stimulation for cervical dystonia. Brain 2007 Nov;130(Pt11):2879-86.

*Kleiner-Fisman G, et al. Subthalamic nucleus deep brain stimulation: summary and meta-analysis of outcomes. Mov Disord 2006 Jun;21(Suppl 14):S290-304.

Kohl S, et al. Deep brain stimulation for treatment-refractory obsessive compulsive disorder: a systematic review. BMC Psychiatry 2014 Aug 2;14:214.

Koy A, et al. Effects of deep brain stimulation in dyskinetic cerebral palsy: a meta-analysis. Mov Disord 2013 May;28(5):747-54.

Koy A, et al. German registry of paediatric deep brain stimulation in patients with childhood-onset dystonia (GEPESTIM). Eur J Paediatr Neurol 2017 Jan;21(1):136-146.

*Krause M, et al. Deep brain stimulation for the treatment of Parkinson's disease: subthalamic nucleus versus globus pallidus internus. J Neurol Neurosurg Psych 2001 Apr;70(4):464-70.

*Krauss JK, et al. Pallidal deep brain stimulation in patients with cervical dystonia and severe cervical dyskinesias with cervical myelopathy. J Neurol Neurosurg Psych 2002 Feb;72(2):249-56.

*Kupsch A, et al. Pallidal deep-brain stimulation in primary generalized or segmental dystonia. NGJM 2006 Nov 9;355(19):1978-90.

*Laxton AW, et al. A phase I trial of deep brain stimulation of memory circuits in Alzheimer's disease. Ann Neurol 2010 Oct;68(4):521-534.

Laxton AW, et al. Deep brain stimulation for the treatment of Alzheimer disease and dementia. World Neurosurg 2013 Sep-Oct;80(3-4):S28.e1-8.

*Lee JY, et al. Thalamic deep brain stimulation for management of essential tremor. J Neurosurg 2005 Sep;103(3):400-3.

*Leone M, et al. Lessons from 8 years' experience of hypothalamic stimulation in cluster headache. Cephalalgia 2008 Jul;28(7):787-97.

*Levy R, et al. Intracranial neurostimulation for pain control: a review. Pain Physician 2010 Mar;13(2):157-65.

Lipsman N, et al. Subcallosal cingulate deep brain stimulation for treatment-refractory anorexia nervosa: a phase 1 pilot study. Lancet 2013 Apr 20;381(9875):1361-70.

*Liu W, et al. Quantitative assessments of the effect of bilateral subthalamic stimulation on multiple aspects of sensorimotor function for patients with Parkinson's disease. Parkinsonism Relat Disord 2005 Dec;11(8):503-8.

Liu Y, et al. Meta-analysis comparing deep brain stimulation of the globus pallidus and subthalamic nucleus to treat advanced Parkinson disease. J Neurosurg 2014 Sep;121(3):709-18.

*Loher TJ, et al. Long-term follow-up study of chronic globus pallidus internus stimulation for post-traumatic hemidystonia. J Neurosurg 2000 Mar;92(3):457-60.

*Loher TJ, et al. Long-term pallidal deep brain stimulation in patients with advanced Parkinson disease: 1-year follow-up study. J Neurosurg 2002 May;96(5):844-53.

SUBJECT: DEEP BRAIN STIMULATION POLICY NUMBER: 7.01.23 CATEGORY: Technology Assessment	EFFECTIVE DATE: 09/16/99 REVISED DATE: 07/19/01, 05/16/02, 03/20/03, 03/18/04, 03/17/05, 01/19/06, 01/18/07, 11/15/07, 11/20/08, 10/29/09, 10/28/10, 09/15/11, 08/16/12, 07/18/13, 06/19/14, 05/29/15, 06/16/16, 05/18/17, 04/19/18 PAGE: 12 OF: 16
---	--

Lozano AM, et al. A multicenter pilot study of subcallosal cingulate area deep brain stimulation for treatment-resistant depression. J Neurosurg 2012 Feb;116(2):315-22.

*Ludwig J, et al. Effects of subthalamic nucleus stimulation and levodopa on the autonomic nervous system in Parkinson's disease. J Neurol Neurosurg Psych 2007 Jul;78(7):742-5.

*Mallet L, et al. Subthalamic nucleus stimulation in severe obsessive-compulsive disorder. NEJM 2008 Nov 13;359(20):2121-34.

*Mayberg H, et al. Deep brain stimulation for treatment-resistant depression. Neuron 2005 Mar 3;45: 651-60.

Mentzel CL, et al. Efficacy and safety of deep brain stimulation in patients with medication-induced tardive dyskinesia and/or dystonia: a systematic review. J Clin Psychiatry 2012 Nov;73(11):1434-8.

Meoni S, et al. Long-term neuropsychiatric outcomes after pallidal stimulation in primary and secondary dystonia. Neurology 2015 Aug 4;85(5):433-40.

Meoni S, et al. Pallidal deep brain stimulation for dystonia: a long term study. J Neurol Neurosurg Psychiatry 2017 Nov;88(11):960-967.

*Merola A, et al. Subthalamic nucleus deep brain stimulation outcome in young onset Parkinson's disease: a role for age at disease onset? J Neurol Neurosurg Psychiatry 2012 Mar;83(3):251-7.

Milev RV, et al. Canadian Network for Mood and Anxiety Treatments (CANMAT) 2016 Clinical; Guidelines for the Management of Adults with Major Depressive Disorder: section 4. neurostimulation treatments. Can J Psychiatry 2016 Sept;61(9):561-575.

Moon W, et al. The cost effectiveness of deep brain stimulation for patients with treatment-resistant obsessive-compulsive disorder. Medicine 2017;96:27.

Morishita T, et al. Deep brain stimulation for treatment-resistant depression: systematic review of clinical outcomes. Neurotherapeutics 2014 Jul;11(3):475-84.

*Moro E, et al. Long-term results of a multicenter study on subthalamic and pallidal stimulation in Parkinson's disease. Mov Disord 2010 Apr 15;25(5):578-86.

Moro E, et al. Efficacy of pallidal stimulation in isolated dystonia: a systematic review and meta-analysis. Eur J Neurol 2017 April;24(4):552-560.

*Mueller J, et al. Pallidal deep brain stimulation improves quality of life in segmental and generalized dystonia: results from a prospective, randomized sham-controlled trial. Mov Disord 2008 Jan;23(1):131-4.

Naesstrom M, et al. A systematic review of psychiatric indications for deep brain stimulation, with focus on major depressive and obsessive-compulsive disorder. Nord J Psychiatry 2016 Oct;70(7):483-491.

*Nakamura K, et al. Effects of unilateral subthalamic and pallidal deep brain stimulation on fine motor functions in Parkinson's disease. Mov Disord 2007 Apr 15;22(5):619-26.

*National Institute for Health and Clinical Excellence. Deep brain stimulation for tremor and dystonia (excluding Parkinson's disease). 2006 Aug [<http://www.nice.org.uk/topic>] accessed 3/13/18.

*National Institute for Health and Clinical Excellence. Deep brain stimulation for Parkinson's disease. 2003 Nov [<http://www.nice.org.uk/topic>] accessed 4/13/17.

National Institute for Health and Clinical Excellence. Deep brain stimulation for intractable trigeminal automatic cephalalgias. 2011 Mar [<http://www.nice.org.uk/topic>] accessed 3/13/18.

SUBJECT: DEEP BRAIN STIMULATION POLICY NUMBER: 7.01.23 CATEGORY: Technology Assessment	EFFECTIVE DATE: 09/16/99 REVISED DATE: 07/19/01, 05/16/02, 03/20/03, 03/18/04, 03/17/05, 01/19/06, 01/18/07, 11/15/07, 11/20/08, 10/29/09, 10/28/10, 09/15/11, 08/16/12, 07/18/13, 06/19/14, 05/29/15, 06/16/16, 05/18/17, 04/19/18 PAGE: 13 OF: 16
---	--

National Institute for Health and Clinical Excellence. Deep brain stimulation for refractory chronic pain syndromes. 2011 Mar [<http://www.nice.org.uk/topic>] accessed 3/13/18.

National Institute for Health and Clinical Excellence. Deep brain stimulation for refractory epilepsy. 2012 Jan [<http://www.nice.org.uk/topic>] accessed 3/13/18.

*Obwegeser AA, et al. Quantitative and qualitative outcome measures after thalamic deep brain stimulation to treat disabling tremors. Neurosurg 2001 Feb;48(2):274-84.

Odekerken VJ, et al. Subthalamic nucleus versus globus pallidus bilateral deep brain stimulation for advanced Parkinson's disease (NSTAPS): a randomized controlled trial. Lancet Neurol 2013 Jan;12(1):37-44.

*Okun MS, et al. Subthalamic deep brain stimulation with a constant-current device in Parkinson's disease: an open-label randomized controlled trial. Lancet Neurol 2012 Feb;11(2):140-9.

Oliveria SF, et al. Safety and efficacy of dual-lead thalamic deep brain stimulation for patients with treatment-refractory multiple sclerosis tremor: a single-center randomized, single-blind, pilot trial. Lancet Neurol 2017 Sept;16(9):691-700.

*Ondo W, et al. Thalamic deep brain stimulation: comparison between unilateral and bilateral placement. Arch Neurol 2001 Feb;58(2):218-22.

Ooms P, et al. Deep brain stimulation for obsessive-compulsive disorders: long-term analysis of quality of life. J Neurol Neurosurg Psychiatry 2014 Feb;85(2):153-8.

*Ostrem JL, et al. Subthalamic nucleus deep brain stimulation in primary cervical dystonia. Neurol 2011 Mar 8;76(10):870-8.

*Owen SL, et al. Deep brain stimulation for the alleviation of post-stroke neuropathic pain. Pain 2006 Jan;120(1-2):202-6.

*Pahwa R, et al. Long-term evaluation of deep brain stimulation of the thalamus. J Neurosurg 2006 Apr;104 (4):506-12.

*Pahwa R, et al. Practice Parameter: treatment of Parkinson disease with motor fluctuations and dyskinesia (an evidence-based review): report of the Quality Standards Subcommittee of the American Academy of Neurology. Neurol 2006 Apr 11;66(7):983-95.

*Pansaon Piedad JC, et al. What patients with Gilles de la Tourette syndrome should be treated with deep brain stimulation and what is the best target? Neurosurg 2012 Jul;71(1):173-92.

Panov F, et al. Deep brain stimulation in DYT1 dystonia: a 10-year experience. Neurosurgery 2013 Jul;73(1):86-93.

Pepper J, et al. Deep brain stimulation versus anterior capsulotomy for obsessive-compulsive disorder: a review of the literature. J Neurosurg 2015 May;122(5):1028-37.

Pereira EA, et al. Thalamic deep brain stimulation for neuropathic pain after amputation or brachial plexus avulsion. Neurosurg Focus 2013 Sep;35(3):E7.

Petrossian MT, et al. Pallidal deep brain stimulation for dystonia: a case series. J Neurosurg Pediatr 2013 Dec;12(6):582-7.

*Piboolnurak P, et al. Levodopa response in long-term bilateral subthalamic stimulation for Parkinson's disease. Mov Disord 2007 May 15;22(7):990-7.

Pollo C, et al. Directional deep brain stimulation: an intraoperative double-blind pilot study. Brain 2014 July;137(Pt 7):2015-2026.

Polo C, et al. Preliminary experience with chronic directional DBS in the STN. Clin Neurophysiol 2016 Sept;127(9):e171.

*Pretto TE, et al. A prospective blinded evaluation of deep brain stimulation for the treatment of secondary dystonia and primary torticollis syndromes. J Neurosurg 2008 Sep;109(3):405-9.

<p>SUBJECT: DEEP BRAIN STIMULATION</p> <p>POLICY NUMBER: 7.01.23</p> <p>CATEGORY: Technology Assessment</p>	<p>EFFECTIVE DATE: 09/16/99</p> <p>REVISED DATE: 07/19/01, 05/16/02, 03/20/03, 03/18/04, 03/17/05, 01/19/06, 01/18/07, 11/15/07, 11/20/08, 10/29/09, 10/28/10, 09/15/11, 08/16/12, 07/18/13, 06/19/14, 05/29/15, 06/16/16, 05/18/17, 04/19/18</p> <p>PAGE: 14 OF: 16</p>
--	---

- *Rasche D, et al. Deep brain stimulation for the treatment of various chronic pain syndromes. Neurosurg Focus 2006 Dec 15;21(6):E8.
- Rebelo P, et al. Thalamic directional deep brain stimulation for tremor: spend less, get more. Brain Stimul 2018 Jan 6. [Epub ahead of print].
- *Rodriguez-Oroz MC, et al. Efficacy of deep brain stimulation of the subthalamic nucleus in Parkinson's disease 4 years after surgery: double blind and open label evaluation. J Neurol Neurosurg Psych 2004 Oct;75(10):1382-5.
- *Rodriguez-Oroz MC, et al. Bilateral deep brain stimulation in Parkinson's disease: a multicentre study with 4 years follow-up. Brain 2005 Oct;128(pt 10):2240-9.
- *Rodrigues JP, et al. Globus pallidus stimulation in advanced Parkinson's disease. J Clin Neurosci 2007 Mar;14(3):208-15.
- *Roh D, et al. Long-term follow-up of deep brain stimulation for refractory obsessive-compulsive disorder. Psychiatry Res 2012 Dec 30;200(2-3):1067-70.
- Rothlind JC, et al. Neuropsychological changes following deep brain stimulation surgery for Parkinson's disease: comparisons of treatment at pallidal and subthalamic targets versus best medical therapy. J Neurol Neurosurg Psychiatry 2015 Jun;86(6):622-29.
- Salanova V, et al. Long-term efficacy and safety of thalamic stimulation for drug-resistant partial epilepsy. Neurology 2015 Mar 10;84(10):1017-25.
- *Schlaepfer TE, et al. Deep brain stimulation for treatment of refractory depression. Lancet 2005 Oct 22-28;366(9495):1420-2.
- Schjerling L, et al. A randomized double-blind crossover trial comparing subthalamic and pallidal deep brain stimulation for dystonia. J Neurosurg 2013 Dec;119(6):1537-45.
- Schrock LE, et al. Tourette syndrome deep brain stimulation: a review and updated recommendations. Mov Disord 2015 Apr;30(4):448-71.
- *Schupbach WM, et al. Neurosurgery at an earlier stage of Parkinson disease: a randomized, controlled trial. Neurol 2007 Jan 23;68(4):267-71.
- Schuepbach WM, et al. Neurostimulation for Parkinson's disease with early motor complications. N Engl J Med 2013 Feb 14;368(7):610-22.
- *Schuurman PR, et al. A comparison of continuous thalamic stimulation and thalamotomy for suppression of severe tremor. NEJM 2000 Feb 17;342(7):505-8.
- Servello D, et al. 16 years of deep brain stimulation for Tourette's syndrome: a critical review. J Neurosurg Sci 2016 June; 60(2):218-229.
- *Sharma A, et al. Efficacy and safety of deep brain stimulation as an adjunct to pharmacotherapy for the treatment of Parkinson disease. Ann Pharmacother 2012 Feb;46(2):248-54.
- *Skogseid IM. Pallidal deep brain stimulation is effective, and improves quality of life in primary segmental and generalized dystonia. Acta Neurol Scand Suppl 2008;188:51-5.
- *Slowinski JL, et al. Unilateral deep brain stimulation of the subthalamic nucleus for Parkinson disease. J Neurosurg 2007 Apr;106(4):626-32.
- Sprengers M, et al. Deep brain and cortical stimulation for epilepsy. Cochrane Database Syst Rev. 2017;6:CD008497.

<p>SUBJECT: DEEP BRAIN STIMULATION</p> <p>POLICY NUMBER: 7.01.23</p> <p>CATEGORY: Technology Assessment</p>	<p>EFFECTIVE DATE: 09/16/99</p> <p>REVISED DATE: 07/19/01, 05/16/02, 03/20/03, 03/18/04, 03/17/05, 01/19/06, 01/18/07, 11/15/07, 11/20/08, 10/29/09, 10/28/10, 09/15/11, 08/16/12, 07/18/13, 06/19/14, 05/29/15, 06/16/16, 05/18/17, 04/19/18</p> <p>PAGE: 15 OF: 16</p>
--	---

Steeves T, et al. Canadian guidelines for the evidence-based treatment of tic disorders: behavioral therapy, deep brain stimulation, and transcranial magnetic stimulation. Can J Psychiatry 2012 Mar;57(3):144-51.

Steigerwald F, et al. Directional deep brain stimulation of the subthalamic nucleus: a pilot study using a novel neurostimulation device. Mov Disord 2016 Aug;31(8):1240-1243.

Taghva AS, et al. Deep brain stimulation for treatment-resistant depression. World Neurosurg 2013 Sep-Oct;80(3-4):S27.e17-24.

Tan ZG, et al. Efficacies of globus pallidus stimulation and subthalamic nucleus stimulation for advanced Parkinson's disease: a meta-analysis of randomized controlled trials. Clin Interv Aging 2016 June 21;11:777-786.

*Temel Y, et al. Single electrode and multiple electrode guided electrical stimulation of the subthalamic nucleus in advanced Parkinson's disease. Neurosurg 2007 Nov;61(5 Suppl 2):346-55.

*Tir M, et al. Exhaustive, one-year follow-up of subthalamic nucleus deep brain stimulation in a large, single-center cohort of Parkinsonian patients. Neurosurg 2007 Aug;61(2):297-304.

*The Deep Brain Stimulation for Parkinson's Disease Study Group. Deep-brain stimulation of the subthalamic nucleus or the pars interna of the globus pallidus in Parkinson's disease. NEJM 2001 Sep;345(35):956-63.

U. S. Food and Drug Administration. Brio neuromodulation system. Summary of safety and effectiveness data. [http://www.accessdata.fda.gov/cdrh_docs/pdf14/P140009b.pdf] accessed 3/13/18.

U. S. Food and Drug Administration. Vercise Deep Brain Stimulation System. Summary of safety and effectiveness data. [https://www.accessdata.fda.gov/cdrh_docs/pdf15/P150031B.pdf] accessed 3/13/18.

Valentin A, et al. Deep brain stimulation of the centromedian thalamic nucleus for the treatment of generalized and frontal epilepsies. Epilepsia 2013 Oct;54(10):1823-33.

*Valdeoriola F, et al. Efficacy and safety of pallidal stimulation in primary dystonia: results of the Spanish multicentric study. J Neurol Neurosurg Psych 2010 Jan;81(1):65-9.

Vercruysse S, et al. Effects of deep brain stimulation of the subthalamic nucleus on freezing of gait in Parkinson's disease: a prospective controlled study. J Neurol Neurosurg Psychiatry 2014 Aug;85(8):871-7.

*Vidailhet M, et al. Bilateral deep-brain stimulation of the globus pallidus in primary generalized dystonia. NEJM 2005 Feb 3;352(5):459-67.

*Vidailhet M, et al. Bilateral, pallidal, deep-brain stimulation in primary generalized dystonia: a prospective 3 year follow-up study. Lancet Neurol 2007 Mar;6(3):201-2.

*Visser-Vanderwalle V, et al. Long-term effects of bilateral subthalamic nucleus stimulation in advanced Parkinson disease: a four year follow-up study. Parkinsonism Relat Disord 2005 May;11(3):157-65.

Voges J, et al. Deep brain stimulation surgery for alcohol addiction. World Neurosurg 2013 Sep-Oct;80(3-4):S28.e21-31.

*Volkman J, et al. Pallidal deep brain stimulation in patients with primary generalized or segmental dystonia: a 5-year follow-up of a randomized trial. Lancet Neurol 2012 Dec;11(12):1029-38.

Volkman J, et al. Pallidal neurostimulation in patients with medication-refractory cervical dystonia: a randomized, sham-controlled trial. Lancet Neurol 2014 Sep;13(9):875-84.

Vonck K, et al. A decade of experience with deep brain stimulation for patients with refractory medial temporal lobe epilepsy. Int J Neural Syst 2013 Feb;23(1):1250034.

Walsh RA, et al. Bilateral pallidal stimulation in cervical dystonia: blinded evidence of benefit beyond 5 years. Brain 2013 Mar;136(Pt3):761-9.

<p>SUBJECT: DEEP BRAIN STIMULATION</p> <p>POLICY NUMBER: 7.01.23</p> <p>CATEGORY: Technology Assessment</p>	<p>EFFECTIVE DATE: 09/16/99</p> <p>REVISED DATE: 07/19/01, 05/16/02, 03/20/03, 03/18/04, 03/17/05, 01/19/06, 01/18/07, 11/15/07, 11/20/08, 10/29/09, 10/28/10, 09/15/11, 08/16/12, 07/18/13, 06/19/14, 05/29/15, 06/16/16, 05/18/17, 04/19/18</p> <p>PAGE: 16 OF: 16</p>
--	---

*Welter ML, et al. Internal pallidal and thalamic stimulation in patients with Tourette's syndrome. Arch Neurol 2008 Jul;65(7):952-7.

Welter ML, et al. Anterior pallidal deep brain stimulation for Tourette's syndrome: a randomized, double-blind, controlled trial. Lancet Neurol 2017 Aug;16(8):610-619.

*Williams A, et al. Deep brain stimulation plus best medical therapy versus best medical therapy alone for advanced Parkinson's disease (PD SURG trial): a randomized, open-label trial. Lancet Neurol 2010 Jun;9(6):581-91.

Wu H, et al. Deep-brain stimulation for anorexia nervosa. World Neurosurg 2013 Sep-Oct;80(3-4):S29.e1-10.

Xie CL, et al. Effects of neurostimulation for advanced Parkinson's disease patients on motor symptoms: a multiple-treatments meta-analysis of randomized controlled trials. Sci Rep May 2016;6:25285.

Xu F, et al. Deep brain stimulation of pallidal versus subthalamic for patients with Parkinson's disease: a meta-analysis of controlled trials. Neuropsychiatr Dis Treat 2016;12:1435-1444.

*Zesiewicz TA, et al. Practice parameter: therapies for essential tremor: report of the Quality Standards Subcommittee of the American Academy of Neurology. Neurol 2005 Jun 28;64(12):2008-20.

*Zesiewicz TA, et al. Evidence-based guideline update: treatment of essential tremor: report of the Quality Standards subcommittee of the American Academy of neurology. Neurology 2011 Nov 8;77(19):1752-5.

Zhang JG, et al. Long-term outcome of globus pallidus internus deep brain stimulation in patients with Tourette syndrome. Mayo Clin Proc 2014 Nov;89(11):1506-14.

Zhang XH, et al. Deep brain stimulation of the globus pallidus internus in patients with intractable Tourette syndrome: a 1-year follow-up study. Chin Med J 2016 May 5;129(9):1022-1027.

Zhang J, et al. The safety issues and hardware-related complications of deep brain stimulation therapy: a single-center retrospective analysis of 478 patients with Parkinson's disease. Clin Interv Aging 2017 June 8;12:923-928.

*Zorzi G, et al. Stimulation of the globus pallidus internus for childhood-onset dystonia. Mov Disord 2005 Sep;20(9):1194-200.

*key article

KEY WORDS:

Brain stimulation, Parkinson's disease, Reclaim, Thalamus, Tremor, dystonia.

CMS COVERAGE FOR MEDICARE PRODUCT MEMBERS

There is currently a National Coverage Determination (NCD) for deep brain stimulation. Please refer to the following NCD website for Medicare Members: <http://www.cms.gov/medicare-coverage-database/details/ncd-details.aspx?NCDId=279&ncdver=1&CoverageSelection=Both&ArticleType=All&PolicyType=Final&s=New+York+-+Upstate&CptHcpcsCode=36514&bc=gAAAABAAAA&>.