

MEDICAL POLICY



MEDICAL POLICY DETAILS	
Medical Policy Title	SMALL BOWEL AND MULTIVISCERAL TRANSPLANTS IN ADULTS AND CHILDREN
Policy Number	7.02.05
Category	Transplants
Effective Date	02/16/00
Revised Date	07/19/01, 06/20/02, 04/24/03, 02/19/04, 02/17/05, 02/16/06, 03/15/07, 03/20/08, 03/19/09, 03/18/10, 03/17/11, 03/15/12, 02/21/13, 02/20/14
Archived Date	02/19/15
Edited Date	03/17/16, 03/16/17, 03/15/18, 03/21/19
Product Disclaimer	<ul style="list-style-type: none"> • If a product excludes coverage for a service, it is not covered, and medical policy criteria do not apply. • If a commercial product (including an Essential Plan product) or a Medicaid product covers a specific service, medical policy criteria apply to the benefit. • If a Medicare product covers a specific service, and there is no national or local Medicare coverage decision for the service, medical policy criteria apply to the benefit.

POLICY STATEMENT

I. Small Bowel Transplant

- A. Based upon our criteria and the review of the peer-reviewed literature, small bowel transplantation has been medically proven to be effective and therefore, **medically appropriate** in pediatric and adult patients with short-bowel syndrome (SBS) for any of the following indications:
1. Impending or overt liver failure due to TPN-induced liver injury. Progressive thrombocytopenia and cholestasis are the most reliable indicators of developing liver dysfunction. Complications of portal hypertension such as variceal bleeding, ascites, and hepatorenal syndrome do not arise until late in the course of disease. Timely referral may allow salvage of the native liver with the more accessible intestinal allograft. Given the higher patient survival rates with this single-organ transplant, patients should be identified and considered for transplant before development of irreversible liver dysfunction.
 2. Thrombosis of 2 or more central veins.
 3. Development of 2 or more episodes of systemic sepsis secondary to line infection per year that requires hospitalization.
 4. A single episode of line-related fungemia, septic shock, and/or acute respiratory distress syndrome.
 5. Frequent episodes of severe dehydration despite intravenous fluid supplementation in addition to TPN.
- B. Based upon our criteria and the lack of peer-reviewed literature, small bowel transplant in adults has not been medically proven to be effective and is considered **investigational** for adults who are able to tolerate TPN.
- C. Based upon our criteria and the lack of peer-reviewed literature, living donations of small bowel for transplantation has not been medically proven to be effective and is considered **investigational**.

II. Multivisceral Transplant

Based upon our criteria and the review of the peer-reviewed literature, multivisceral transplantation has been medically proven to be effective and therefore, **medically appropriate** in pediatric and adult patients with intestinal failure and concurrent liver failure.

III. Candidates must meet all of the following criteria:

- A. Adequate cardiopulmonary status;
- B. Absence of active infection;

Medical Policy: SMALL BOWEL AND MULTIVISCERAL TRANSPLANTS IN ADULTS AND CHILDREN

Policy Number: 7.02.05

Page: 2 of 5

- C. Absence of malignancy (other than non-melanoma skin cancers), or malignancy has been completely resected, or (upon medical review) it is determined that malignancy has been treated with small likelihood of recurrence and acceptable future risks; and
 - D. Documentation of patient compliance with medical management.
- IV. Transplant is contraindicated in patients with HIV infection unless *ALL* of the following criteria are met:
- A. CD4 count greater than 200 cells/mm³,
 - B. HIV-1RNA undetectable,
 - C. On stable anti-retroviral therapy greater than 3 months, and
 - D. No other complications from AIDS (e.g., opportunistic infection, including aspergillus, tuberculosis, coccidioidomycosis; resistant fungal infections, Kaposi's sarcoma, or other neoplasm).

POLICY GUIDELINES

- I. Prior authorization is contract dependent. Approvals for all transplants, including arrangements with an approved transplant center, may be required.
- II. Pre-transplant evaluation documentation could include the following clinical information. If testing is unable to be performed, the rationale for not performing the testing should be included in the documentation:
 - A. Clinical Evaluation:
 - 1. Confirmation of diagnosis;
 - 2. Identification of comorbidities;
 - 3. Treatment of co-morbidities;
 - 4. Current assessment of co-morbidities;
 - 5. Consult notes (if applicable).
 - B. Psycho-Social Evaluation:
 - 1. Karnofsky performance score;
 - 2. Identification of stressors (family support, noncompliance issues, motivational issues, alcohol or substance abuse).
 - C. Dental Evaluation.
 - D. Lab Tests:
 - 1. CBC, metabolic profile;
 - 2. Serologies: CMV,
 - 3. Hepatitis B and C;
 - 4. HIV Testing.
 - E. Cardiac Assessment:
 - 1. 12Lead EKG;
 - 2. Stress echo or MUGA Scan.
 - F. Pulmonary Assessment:
 - 1. Chest x-ray;
 - 2. Pulmonary function tests (PFTs);
 - 3. Low dose screening CT for individuals considered high-risk for lung cancer (e.g., 20-30 pack history of smoking).
 - G. Age Appropriate Screening Tests:
 - 1. Age greater than or equal to 50 years:
 - a. Colonoscopy (within 10 years); or
 - b. Flexible sigmoidoscopy (within 5 years); or
 - c. Guaiac stool testing (within 1 year); or
 - d. Rationale of contraindication to testing (if applicable).
 - 2. Women age 21-70 years:

Medical Policy: SMALL BOWEL AND MULTIVISCERAL TRANSPLANTS IN ADULTS AND CHILDREN

Policy Number: 7.02.05

Page: 3 of 5

- a. Pap smear (within 3 years).
3. Women age greater than or equal to 40 years:
 - a. Mammogram (within 2 years).

DESCRIPTION

Small Bowel Transplant

The purpose of a small bowel (SB) transplant is to restore bowel function and allow for adequate nutrition in patients with short bowel syndrome (SBS). It may be an alternative to total parenteral nutrition (TPN) for selected patients who are predicted to have poor survival on TPN.

Multivisceral Transplant

Candidates for multivisceral (MV) transplant have short bowel syndrome and terminal liver failure or other gastrointestinal problems such as pancreatic failure, thromboses of the celiac axis and the mesenteric artery or pseudo-obstruction affecting the entire gastrointestinal tract. Due to anatomic or other medical problems this group of patients requires a more extensive transplant procedure than a small bowel and liver. MV transplantation may include in addition to the small bowel and liver, stomach, duodenum, jejunum, ileum, pancreas and/or colon.

MV transplantation is an infrequently performed procedure, but without this procedure most patients face 100% mortality.

RATIONALE

Total parenteral nutrition is the only established treatment that can produce long-term survival once the small intestine is dysfunctional and oral nutrition is ineffective. TPN requires placement of a permanent venous access device. There are some serious, life threatening complications that can occur as a result of TPN including hepatobiliary disease, thrombosis due to the venous catheter, or sepsis from the venous access line.

There are limited long-term data on small bowel and multivisceral transplants due to the small numbers performed. International Intestinal Transplant Registry outcomes published in 1999 include overall patient and graft survival rates of 69% for isolated intestine recipients at 1 year and 66% and 63% for liver/bowel and multivisceral grafts. It is possible that some patient with severe TPN-associated complications face a higher probability of mortality with continued medical management than the risk of transplantation. Small bowel and multivisceral transplantation is reserved for selected patients with life threatening complications of TPN.

Living donor isolated or combined liver/intestinal transplants have been studied in very small case studies. Typically living donor transplants have been reserved for children who are at high risk for premature death while on the cadaveric waiting list and have no central venous access or for children with impending TPN-related liver failure. A living donor liver transplant may be performed first followed by an intestinal transplant from the same donor later. Advantages to living donors transplants include better HLA matching, reduction of cold ischemia time, and no wait-listing for a transplant, thus the patient is less likely to die while waiting for an organ. Results from the studies show little or no complications after transplant for the donor. Most complications such as diarrhea, weight loss, and nausea are resolved within a few weeks of surgery. However these small studies are lacking long-term follow up of the donors. Patient survival and graft survival for recipients of living donor combined liver/intestinal or isolated intestinal transplants has been favorable. More large studies are needed to determine if patient survival is comparable or better than those patients receiving cadaveric organs. Most studies suggest that living donor transplanted organs are to be reserved for those circumstances where there is high risk for death and no cadaveric donors are available.

Solid organ transplantation for candidates that are HIV positive has long been controversial, due to the long-term prognosis for HIV positivity, and the impact of immunosuppression on HIV disease. Although HIV+ transplant recipients may be a research interest of some transplant centers, the minimal data regarding long-term outcome in these patients consist primarily of case reports and abstract presentations of liver and kidney recipients. Nevertheless, some transplant surgeons would argue that HIV positivity is no longer an absolute contraindication to transplant due to the advent of highly active antiretroviral therapy (HAART), which has markedly changed the natural history of the disease. Furthermore, UNOS states that asymptomatic HIV+ patients should not necessarily be excluded for candidacy for organ transplantation, stating "A potential candidate for organ transplantation whose test for HIV is positive but who is in an

Medical Policy: SMALL BOWEL AND MULTIVISCERAL TRANSPLANTS IN ADULTS AND CHILDREN

Policy Number: 7.02.05

Page: 4 of 5

asymptomatic state should not necessarily be excluded from candidacy for organ transplantation, but should be advised that he or she may be at increased risk of morbidity and mortality because of immunosuppressive therapy". In 2001, the Clinical Practice Committee of the American Society of Transplantation proposed that the presence of AIDS could be considered a contraindication to small bowel and multivisceral transplants unless the specific criteria were present. These criteria are listed in this policy regarding HIV status and small bowel and multivisceral transplants.

CODES

- Eligibility for reimbursement is based upon the benefits set forth in the member's subscriber contract.
- CODES MAY NOT BE COVERED UNDER ALL CIRCUMSTANCES. PLEASE READ THE POLICY AND GUIDELINES STATEMENTS CAREFULLY.
- Codes may not be all inclusive as the AMA and CMS code updates may occur more frequently than policy updates.

CPT Codes

Code	Description
44120	Enterectomy, resection of small intestine; single resection and anastomosis
44121	;each additional resection and anastomosis (List separately in addition to code for primary procedure)
44125	with enterostomy
44135	Intestinal allotransplantation from a cadaver donor
44136 (E/I)	Intestinal allotransplantation from a living donor
44137	Removal of transplanted intestinal allograft, complete
47135	Liver allotransplantation; orthotopic, partial or whole from cadaver or living donor, any age

Copyright © 2019 American Medical Association, Chicago, IL

HCPCS Codes

Code	Description
S2053	Transplantation of small intestine, and liver allografts
S2054	Transplantation of multivisceral organs

ICD10 Codes

Code	Description
K72.10	Chronic hepatic failure without coma
K72.11	Chronic hepatic failure with coma
K72.90	Hepatic failure, unspecified without coma
K72.91	Hepatic failure, unspecified with coma
K91.2	Postsurgical malabsorption, not elsewhere classified

REFERENCES

*Benedetti E, et al. Progressive functional adaptation of segmental bowel graft from living related donor. Transplant 2001;71(4):569-71.

Beyer-Berjot L, et al. Intestinal transplantation: indications and prospects. J Visc Surg 2012 Dec;149(6):380-4.

*BlueCross BlueShield Technology Assessment Program. Small bowel transplants in adults and multivisceral transplants in adults and children. 1999 Jul;14(9).

BlueCross BlueShield Association. Medical Policy Reference Manual Policy #7.03.04. Isolated small bowel transplant. 2017 Aug 08.

Proprietary Information of Excellus Health Plan, Inc.

Medical Policy: SMALL BOWEL AND MULTIVISCERAL TRANSPLANTS IN ADULTS AND CHILDREN

Policy Number: 7.02.05

Page: 5 of 5

BlueCross BlueShield Association. Medical Policy Reference Manual Policy #7.03.05. Small bowel/liver and multivisceral transplant. 2017 Aug 10.

Farmer DG, et al. Pretransplant predictors of survival after intestinal transplantation: analysis of a single-center experience of more than 100 transplants. Transplantation 2010 Dec 27;90(12):1574-80.

*Gangemi A, et al. Lessons learned in pediatric small bowel and liver transplantation from living-related donors. Transplantation 2009;87:1027-30.

Gotthardt DN, et al. Indications for intestinal transplantation: recognizing the scope and limits of total parenteral nutrition. Clin Transplant 2013;27(S25):49-55.

Grant D, et al. Intestinal transplant registry report: global activity and trends. Am J Transplant 2015 Jan;15(1):210-9.

*Horslen SP, et al. Isolated liver transplantation in infants with end-stage liver disease associated with short bowel syndrome. Ann Surg 2002 Mar;235(3):435-9.

Mangus RS, et al. Multivisceral transplantation: expanding indications and improving outcomes. J Gastrointest Surg 2013 Jan;17(1):179-87.

Middleton SJ, et al. Adult small intestinal and multivisceral transplantation: lessons through the “retrospecto-scope” at a single UK centre from 1991 to 2013. Transplant Proc 2014 Jul-Aug;46(6):2114-8.

Nayyar N, et al. Pediatric small bowel transplantation. Semin Pediatr Surg 2010 Feb;19(1):68-77.

*Sokal EM, et al. Liver and intestinal transplantation in children: working group report of the first world congress of pediatric gastroenterology, hepatology, and nutrition. J Ped Gastroenterol Nutrit 2002 Aug;35:S159-72.

Sudan D. The current state of intestine transplantation: indications, techniques, outcomes and challenges. Am J Transplant 2014 Sep;14(9):1976-84.

Trevizol AP, et al. Intestinal and multivisceral retransplantation results: literature review. Transplant Proc 2013 Apr;45(3):1133-6.

Ueno T, et al. Current status of intestinal transplantation. Surg Today 2010;40:1112-22.

Varkey J, et al. Survival of patients evaluated for intestinal and multivisceral transplantation - the Scandinavian experience. Scand J Gastroenterol 2013;48(6):702-11.

*Key Article

KEY WORDS

Intestine, Multivisceral Small bowel, Transplant

CMS COVERAGE FOR MEDICARE PRODUCT MEMBERS

There is currently a National Coverage Determination (NCD) for Intestinal and Multi-Visceral Transplantation. Please refer to the following NCD website for Medicare Members: http://www.cms.gov/medicare-coverage-database/details/ncd-details.aspx?NCDId=280&ncd_ver=2&bc=BAABAAAAAAAA&