

MEDICAL POLICY



MEDICAL POLICY DETAILS	
Medical Policy Title	MOLECULAR MARKERS IN FINE NEEDLE ASPIRATES OF THE THYROID
Policy Number	2.02.49
Category	Laboratory Test
Effective Date	11/19/15
Revised Date	09/15/16, 09/21/17, 12/20/18
Product Disclaimer	<ul style="list-style-type: none"> • If a product excludes coverage for a service, it is not covered, and medical policy criteria do not apply. • If a commercial product (including an Essential Plan product) or a Medicaid product covers a specific service, medical policy criteria apply to the benefit. • If a Medicare product covers a specific service, and there is no national or local Medicare coverage decision for the service, medical policy criteria apply to the benefit.

POLICY STATEMENT

- I. Based on our criteria and assessment of the peer-reviewed literature, mutation analysis in fine-needle aspirates of the thyroid (e.g., ThyroSeq[®], ThyGenX Thyroid Oncogene Panel) is considered investigational.
- II. Based on our criteria and assessment of the peer-reviewed literature, the use of a gene expression classifier in fine-needle aspirates of the thyroid (e.g., Afirma[®] Gene Expression Classifier) that are cytologically considered to be indeterminate, atypical or suspicious for malignancy is considered **medically appropriate** when *surgical decisions will be based on test results* for thyroid nodules with the following:
 - A. Cytological diagnosis of atypia of undetermined significance/follicular lesion of either undetermined significance (AUS/FLUS) on fine-needle aspiration (FNA); OR
 - B. Cytological diagnosis of follicular neoplasm/suspicious for follicular neoplasm (FN/SFN) on fine-needle aspiration (FNA); OR
 - C. Bethesda III or IV on FNA cytology (please refer to the Bethesda criteria in the Description section); and
 - D. Size greater than 1.0 cm; and
 - E. Without clinical suspicion of malignancy based on provider judgment and ultrasonography; and
 - F. No compressive manifestations.
- III. Based on our criteria and assessment of the peer-reviewed literature, the use of a gene expression classifier in fine-needle aspirates of the thyroid not meeting the above criteria are considered investigational.

POLICY GUIDELINES

- I. Ultrasound features associated with low suspicion of malignancy include:
 - A. Isoechoic or hyperechoic solid nodules without microcalcifications; or
 - B. Mixed solid/cystic nodules without microcalcification; or
 - C. Spongiform nodules.
- II. The ThyroSeq[®] v.2 Next Generation Sequencing (NGS) panel (CBLPath, Ocala, FL) includes sequencing of more than 60 genes and per manufacturer's website is indicated when FNA cytology indicates atypical of uncertain significance (AUS) or follicular lesion of undetermined significance (FLUS), follicular neoplasm (FN) or suspicious for follicular neoplasm, or suspicious for malignancy.
- III. The ThyraMIR Micro RNA[™]/ThyGenX[®] Thyroid Oncogene Panel (Interpace Diagnostics, Parsippany, NJ) is a Next Generation Sequencing panel designed to be used in patients with indeterminate thyroid FNA results. The panel includes sequencing of 8 genes associated with papillary thyroid carcinoma (PTC) and follicular carcinomas. The ThyraMIR[™] is a miRNA gene expression classifier and is based on the evaluation of expression of 10 miRNAs.

Medical Policy: MOLECULAR MARKERS IN FINE NEEDLE ASPIRATES OF THE THYROID

Policy Number: 2.02.49

Page: 2 of 6

When used in combination ThyraMIR™ can identify malignancy in nodules that are negative for ThyGenX® which potentially improves overall sensitivity and ability to detect malignancy.

- IV. The Afirma® Gene Expression Classifier (Afirma GEC; Veracyte, San Francisco, CA) analyzes the expression of 142 different genes to determine patterns associated with benign finding on surgical biopsy. It is indicated for thyroid nodules that have an indeterminate classification on FNA. Testing is limited to 2 thyroid nodules.
- V. The Afirma® BRAF and Afirma MTC are two tests which use mRNA expression-based classification to evaluate for BRAF mutations or mutations associated with medullary thyroid carcinoma. Both tests are options when the Afirma GEC results are malignant or suspicious.
- VI. The Federal Employee Health Benefit Program (FEHBP/FEP) requires that procedures, devices or laboratory tests approved by the U.S. Food and Drug Administration (FDA) may not be considered investigational and thus these procedures, devices or laboratory tests may be assessed only on the basis of their medical necessity.

DESCRIPTION

Thyroid nodules are common endocrine tumors which appear as palpable nodules in 5% of adults. Up to 50% of women and 20% of men over age 50 on ultrasound or autopsy studies are found to have thyroid nodules. The majority of thyroid nodules are benign but 10-15% may be malignant, most often as papillary thyroid cancer. Fine needle aspiration (FNA) is performed in nodules which require biopsy with sufficient information obtained to classify the majority of nodules as benign and a smaller percentage as malignant. However 15-30% of aspirations yield indeterminate cytology including subtypes such as atypia of undetermined significance (AUS) or follicular lesion of undetermined significance (FLUS), suspicious for follicular neoplasm (FN) or suspicious for malignancy. Current guidelines recommend either partial (lobectomy) or complete thyroidectomy for those nodules determined as malignant and those of indeterminate cytology. On histological evaluation only 15-30% of the thyroid nodules of indeterminate cytology are malignant consequently many patients undergo surgery for benign disease when expectant management or other treatments would have been more appropriate with a retrospective assessment. Due to the limitations of the FNA, other methods to assist in determining whether a nodule is benign or malignant prior to surgery have been developed. The ThyroSeq and ThyGenX thyroid Oncogene panel are two tests evaluation for point mutation associated with thyroid cancers using next-generation sequencing. The Afirma Gene Expression Classifier (GEC) analyzes genetic alterations through the use of gene expression profiling.

RATIONALE

Analysis for mutations associated with thyroid cancer in fine needle aspirates (FNA) of the thyroid that are cytologically indeterminate has a high positive predictive value for malignancy. However, patients with an equivocal FNA result would likely proceed to surgery regardless of mutation status, with intraoperative consultation to guide the necessity and extent of surgery. Mutation analysis has the potential to improve the accuracy of an equivocal FNA of the thyroid and may play a role in preoperative risk stratification and surgical planning. Studies suggest that testing for a panel of mutations associated with thyroid cancer may allow the appropriate selection of patients for surgical management with an initial total thyroidectomy. Additional studies are needed to validate these results. Mutation analysis does not achieve a high enough negative predictive value (NPV) to identify which patients can undergo watchful waiting over thyroid surgery. Although the presence of certain mutations may predict more aggressive malignancies, the clinical utility of identifying these mutations preoperatively has not been established.

There is one commercially available gene expression classifier (GEC) that has been developed to predict benignancy in thyroid nodules. The reported NPV of the GEC in predicting which thyroid nodules with indeterminate cytology are benign is high. Studies in which patients who avoided surgery based on GEC results need longer follow-up data.

Medical Policy: MOLECULAR MARKERS IN FINE NEEDLE ASPIRATES OF THE THYROID

Policy Number: 2.02.49

Page: 3 of 6

The American Thyroid Association (ATA) statement on Surgical Application of Molecular Profiling for Thyroid Nodules (2015) states techniques for molecular profiling of thyroid cytology specimens have evolved as adjuncts to guide the appropriate management of cytologically indeterminate nodules. However, it must be stressed that the utility of any molecular test is only applicable clinically when combined with clinical and sonographic risk factors for malignancy and with understanding of the prevalence of malignancy for the Bethesda cytologic categories at the reporting institution. Future studies on further refinements and expansion of gene sets in analytic panels will likely improve the diagnostic accuracy of molecular analyses of thyroid cytology specimens and offer promise for personalizing surgical therapy, with the potential for cost and risk reduction in the diagnostic and therapeutic approaches to treating differentiated thyroid cancer.

The National Comprehensive Cancer Network (NCCN) Guidelines for Thyroid Carcinoma (2017) state that the choice of the precise molecular test depends on the cytology and the clinical question being asked. Indeterminate groups include follicular or Hürthle cell neoplasms and atypia of undetermined significance (AUS) or follicular lesion of undetermined significance (FLUS). The NCCN panel recommends molecular diagnostic testing for evaluating FNA results that are suspicious for follicular cell neoplasms or AUS/FLUS that are low clinical suspicion of malignancy. Molecular diagnostic testing is not recommended for suspected Hürthle cell neoplasms. Molecular diagnostic testing may include multigene assays (e.g., the gene expression classifier) or individual mutational analysis. Patients can be followed with observation if the application of a specific molecular diagnostic test (in conjunction with clinical and ultrasound features) results in a predicted risk of malignancy that is comparable to the rate seen in cytologically benign fine needle aspiration (FNA). It is important to note that the positive predictive value of molecular diagnostics may be significantly influence by the pre-test probability of disease associated with the various FNA cytology groups. Furthermore, in the cytologically indeterminate groups, the risk of malignancy for FNA can vary widely between institutions. Proper implementation of molecular diagnostics into clinical care requires an understanding of both the performance characteristics of the molecular test and its clinical meaning across a range of pre-test probabilities.

In 2007, the National Cancer Institute (NCI) Thyroid FNA State of Science Conference, developed the *Bethesda System for Reporting Thyroid Cytopathology*. The purpose of the conference was to develop a uniform reporting system for thyroid FNA to facilitate effective communication among cytopathologist, endocrinologists, surgeons, radiologists, and other health care providers.

The Bethesda System for Reporting Thyroid Cytopathology

Risk Category	Definition	Diagnostics
I	Nondiagnostic or Unsatisfactory	Cyst fluid only Virtually acellular specimen Other (obscuring blood, clotting artifact, etc.)
II	Benign	Consistent with a benign follicular nodule (includes adenomatoid nodule, colloid nodule, etc.) Consistent with lymphocytic (Hashimoto) thyroiditis in the proper clinical context Consistent with granulomatous (subacute) thyroiditis Other
III	Atypia of Undetermined Significance <i>or</i> Follicular Lesion of Undetermined Significance	
IV	Follicular Neoplasm or Suspicious for a Follicular Neoplasm	Specify if Hürthle cell (oncocytic) type

Medical Policy: MOLECULAR MARKERS IN FINE NEEDLE ASPIRATES OF THE THYROID**Policy Number: 2.02.49****Page: 4 of 6**

V	Suspicious for Malignancy	Suspicious for papillary carcinoma Suspicious for medullary carcinoma Suspicious for metastatic carcinoma Suspicious for lymphoma Other
VI	Malignant	Papillary thyroid carcinoma Poorly differentiated carcinoma Medullary thyroid carcinoma Undifferentiated (anaplastic) carcinoma Squamous cell carcinoma Carcinoma with mixed features (specify) Metastatic carcinoma Non-Hodgkin lymphoma Other

CODES

- Eligibility for reimbursement is based upon the benefits set forth in the member's subscriber contract.
- **CODES MAY NOT BE COVERED UNDER ALL CIRCUMSTANCES. PLEASE READ THE POLICY AND GUIDELINES STATEMENTS CAREFULLY.**
- Codes may not be all inclusive as the AMA and CMS code updates may occur more frequently than policy updates.

CPT Codes

Code	Description
81445	Targeted genomic sequence analysis panel, solid organ neoplasm, DNA analysis, and RNA analysis when performed, 5-50 genes (eg, ALK, BRAF, CDKN2A, EGFR, ERBB2, KIT, KRAS, NRAS, MET, PDGFRA, PDGFRB, PGR, PIK3CA, PTEN, RET), interrogation for sequence variants and copy number variants or rearrangements, if performed
81479	Unlisted molecular pathology procedure
81545	Oncology (thyroid), gene expression analysis of 142 genes, utilizing fine needle aspirate, algorithm reported as a categorical result (e.g., benign or suspicious)
0018U	Oncology (thyroid), microRNA profiling by RT-PCR of 10 microRNA sequences, utilizing fine needle aspirate, algorithm reported as a positive or negative result for moderate to high risk of malignancy (<i>ThyraMIR™</i> , <i>Interpace Diagnostics</i>)
0026U	ONC THYR DNA&MRNA 112 GENES (<i>Thyroseq Genomic Classifier™</i>)

Copyright © 2019 American Medical Association, Chicago, IL

HCPCS Codes

Code	Description
No specific codes	

ICD10 Codes

Code	Description
C73	Malignant neoplasm of thyroid gland
D34	Benign neoplasm of thyroid gland
D44.0	Neoplasm of uncertain behavior of thyroid gland

Proprietary Information of Excellus Health Plan, Inc.

Medical Policy: MOLECULAR MARKERS IN FINE NEEDLE ASPIRATES OF THE THYROID

Policy Number: 2.02.49

Page: 5 of 6

REFERENCES

- *Alexander EK, et al. Preoperative diagnosis of benign thyroid nodules with indeterminate cytology. N Engl J Med 2012 Aug 23;267(8):705-15.
- Al-Qurayshi Z, et al. Association of malignancy prevalence with test properties and performance of the gene expression classifier in indeterminate thyroid nodules. JAMA Otolaryngol Head Neck Surg 2017 Apr 1;143(4):403-408.
- BlueCross BlueShield Association. Molecular markers in fine needle aspirates of the thyroid. Medical Policy Reference Manual Policy #2.04.78. 2016 Nov 10.
- *Cibas ES, et al. The Bethesda System For Reporting Thyroid Cytopathology. Am J Clin Pathol 2009 Nov;132(5):658-65.
- Gharib H, et al. American Association of Clinical Endocrinologists, American College of Endocrinology, and Associazione Medici Endocrinologi Medical Guidelines for Clinical Practice for the Diagnosis and Management of Thyroid Nodules--2016 update. Endocr Pract. 2016 May;22(5):622-39.
- Harrison G, et al. Evaluation of the Afirma gene expression classifier in repeat indeterminate thyroid nodules. Arch Pathol Lab Med 2017 Jul;141(7):985-989.
- Jin H, et al. BRAFV600E mutation as a predictor of thyroid malignancy in indeterminate nodules: a systematic review and meta-analysis. Eur J Surg Oncol 2017 Jul;43(7):1219-1227.
- Jug RC, et al. Molecular testing for indeterminate thyroid nodules: performance of the Afirma Gene Expression Classifier and ThyroSeq panel. Cancer Cytopathol. 2018 Apr 10 [Epub ahead of print].
- Kay-Rivest E, et al., The first Canadian experience with the Afirma[®] gene expression classifier test. J Otolaryngol Head Neck Surg 2017 Apr 4;46(1):25.
- Livhits MJ, et al. Gene expression classifier vs targeted next-generation sequencing in the management of indeterminate thyroid nodules. J Clin Endocrinol Metab 2018;103(6):2261-2264.
- National Comprehensive Cancer Network (NCCN). Clinical practice guidelines in oncology. Thyroid carcinoma. V.2.2017. [https://www.nccn.org/professionals/physician_gls/pdf/thyroid.pdf] accessed 09/19/18 03:17.
- *Nikiforov YE, et al. Highly accurate diagnosis of cancer in thyroid nodules with follicular neoplasm/suspicious for a follicular neoplasm cytology by ThyroSeq v2 next- generation sequencing assay. Cancer 2014 Dec 1;120(23):3627-34.
- Sacks WL, et al. Impact of Afirma gene expression classifier on cytopathology diagnosis and rate of thyroidectomy. Cancer Cytopathol 2016 Oct;124(10):722-728.
- Taye A, et al. Clinical performance of a next-generation sequencing assay (ThyroSeq vs) in the evaluation of indeterminate thyroid nodules. Surgery. 2018 Jan;163(1):97-103.
- Wu JX, et al. Effect of malignancy rates on cost-effectiveness of routine gene expression classifier testing for indeterminate thyroid nodules. Thyroid 2016 Jul;26(7):916-22.
- Yip L, Sossa JA. Molecular-directed treatment of differentiated thyroid cancer – advances in diagnosis and treatment. JAMA Surg 2016 Jul 1;151(7):663-70.

*Key Article

KEY WORDS

ThyroSeq[®], Afirma[®] Gene Expression Classifier (GEC), fine needle aspiration of the thyroid, molecular markers of thyroid, ThyGenX Thyroid Oncogene Panel

Medical Policy: MOLECULAR MARKERS IN FINE NEEDLE ASPIRATES OF THE THYROID

Policy Number: 2.02.49

Page: 6 of 6

CMS COVERAGE FOR MEDICARE PRODUCT MEMBERS

There is currently a Local Coverage Determination (LCD) for Molecular Pathology Procedures. Please refer to the following LCD website for Medicare members: https://www.cms.gov/medicare-coverage-database/details/lcd-details.aspx?LCDId=35000&ver=128&CntrctrSelected=298*1&Cntrctr=298&s=41&DocType=Active&bc=AggAAAIBAAAA&

There is currently a Local Coverage Article (LCA) for Molecular Pathology Procedures-Related to Molecular Policy Procedures LCD (L35000) (A56199). Please refer to the following LCD website for Medicare members: https://www.cms.gov/medicare-coverage-database/details/article-details.aspx?articleId=56199&ver=11&LCDId=35000&CntrctrSelected=298*1&Cntrctr=298&s=41&DocType=Active&bc=AggAAAIBAAAA&